

Multitensors and monads on categories of enriched graphs

Mark Weber

ABSTRACT. In this paper we unify the developments of [3], [6] and [11] into a single framework in which the interplay between multitensors on a category V , and monads on the category $\mathcal{GV}$ of graphs enriched in V , is taken as fundamental. The material presented here is the conceptual background for subsequent work: in [5] the Gray tensor product of 2-categories and the Crans tensor product [13] of Gray categories are exhibited as existing within our framework, and in [27] the explicit construction of the funny tensor product of categories is generalised to a large class of Batanin operads.

CONTENTS

1. Introduction	1
2. Connected objects and locally c-presentable categories	4
3. Categories of enriched graphs	11
4. The multitensor to monad construction	20
5. The 2-functoriality of the multitensor to monad construction	33
6. Distributive laws and monoidal monads	43
7. Contractible operads and multitensors	53
8. Trimble's construction	56
Acknowledgements	59
References	59

1. Introduction

In [3] the problem of how to give an explicit combinatorial definition of weak higher categories was solved, and the development of a conceptual framework for their further analysis was begun. In the aftermath of this, the expository work of other authors, most notably Street [25] and Leinster [21], contributed greatly to our understanding of these ideas. The central idea of [3] is that the description of any n -dimensional categorical structure X , may begin by starting with just the underlying n -globular set, that is, the sets and functions

$$X_0 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_1 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_2 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_3 \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} \cdots \begin{array}{c} \xleftarrow{s} \\ \xleftarrow{t} \end{array} X_n$$

1

satisfying the equations $ss = st$ and $ts = tt$, which embody the the objects (elements of X_0), arrows (elements of X_1) and higher cells of the structure in question. At this stage no compositions have been defined, and when they are, one has a globular set with extra structure. In this way the problem of defining an n -categorical structure of a given type is that of defining the monad on the category $\widehat{\mathbb{G}}_{\leq n}$ of n -globular sets whose algebras are these structures.

As explained in the introduction to [6], this approach works because the monads concerned have excellent formal properties, which facilitate their explicit description and further analysis. The n -operads of [3] can be defined from the point of view of monads: one has the monad $\mathcal{T}_{\leq n}$ on $\widehat{\mathbb{G}}_{\leq n}$ whose algebras are strict n -categories, and an n -operad consists of another monad A on $\widehat{\mathbb{G}}_{\leq n}$ equipped with a cartesian monad morphism $A \rightarrow \mathcal{T}_{\leq n}$. The algebras of this n -operad are just the algebras of A .

Strict n -categories are easily defined by iterated enrichment: a strict $(n+1)$ -category is a category enriched in the category of strict n -categories via its cartesian product, but are too strict for the intended applications in homotopy theory and geometry. For $n = 3$ the strictest structure one can replace an arbitrary weak 3-category with – and not lose vital information – is a Gray category, which is a category enriched in 2-Cat using the Gray tensor product of 2-categories instead of its cartesian product [16]. This experience leads naturally to the idea of trying to define what the higher dimensional analogues of the Gray tensor product are, so as to set up a similar inductive definition as for strict n -categories, but to capture the appropriate semi-strict n -categories, which in the appropriate sense, would form the strictest structure one can replace an arbitrary weak n -category with and not lose vital information. The search for such an inductive definition is the main motivation for this series of papers which began with [6]. This article is the next in the series, and is itself followed by [5] and [27]. Note also that this article and [5] taken together amount to an elaboration of the contents of [4].

A multitensor on a category V is exactly a lax monoidal structure on V . In [6] a description of the relationship between multitensors and the higher operads of [3] was begun. Sections(4) and (5) of the present paper continue this development by studying, for a given category V , the passage

$$\text{Multitensors on } V \mapsto \text{Monads on } \mathcal{GV}$$

where $\mathcal{GV}$ is the category of graphs enriched in V , in a systematic way in which the above assignment is seen as the object part of a 2-functor

$$\Gamma : \text{DISTMULT} \rightarrow \text{MND}(\mathbf{CAT}/\mathbf{Set})$$

where DISTMULT is a sub 2-category of the 2-category of lax monoidal categories, and MND is as defined by the formal theory of monads [24]. From this perspective, one is able to describe in a more efficient and general way, many of the previous developments of higher category theory in the globular style. For instance, in section(6.3) we give a short and direct explicit construction of the monads $\mathcal{T}_{\leq n}$ for strict n -categories from which all their key properties are easily witnessed. Throughout sections(4) and (5) we give shorter and more general proofs of some of the main results of [6]. In section(8) we apply our theory to give an efficient general account of Cheng's analysis [11] of the Trimble definition of weak n -category.

It is an important observation of Street [24] that monads in $\text{MND}(\mathcal{K})$, where $\mathcal{K}$ is an arbitrary 2-category, are exactly distributive laws in the sense of Beck [7].

This fact and our expression of Γ as a 2-functor explains the ubiquity of monad distributive laws in this subject – see [10] [11]. Moreover in section(6) we discuss the sense in which a multitensor and a monad can distribute, leading to an analogous theory to that of [7], in which the role of monad distributive laws is played by the coherence data for a monoidal or opmonoidal monad.

The central construction of a monad from a multitensor gets along very well with existing categorical notions. The main result expressing this is theorem(29). In preparation for this result we review in section(2) some of the relevant theory of connected objects within lexensive categories and the notion of a locally c-presentable category – which is a mild strengthening of the well-known notion of locally presentable category. This notion is important for us since categories of algebras of higher operads are locally c-presentable. Then in section(3) we give a systematic account of how the passage $V \mapsto \mathcal{G}V$ respects various categorical properties.

A weak n -category is an algebra of a *contractible* n -operad¹. In section(7) we recall this notion, give an analogous notion of contractible multitensor, and in corollary(59), give the canonical relationship between them.

In this paper we operate at a more abstract level than in much of the previous work on this subject. In particular, instead of studying monads on the category of n -globular sets, or even on presheaf categories, we work with monads defined on some category $\mathcal{G}V$ of enriched graphs. As our work shows, the main results and notions of higher category theory in the globular style can be given in this setting. So one could from the very beginning start not with **Set** as the category of 0-categories, but with a nice enough V . For all the constructions to go through, such as that of $\mathcal{T}_{\leq n}$, the correspondence between monads and lax tensor products, their lifting theorem, as well as the very definition of weak n -category, it suffices to take V to be a locally c-presentable category.

Proceeding this way one obtains then the theory of n -dimensional structures enriched in V . That is to say, the object of n -cells between any two $(n-1)$ -cells of such a structure would be an object of V rather than a mere set. Some alternative choices of V which could perhaps be of interest are: (1) the ordinal $[1] = \{0 < 1\}$ (for the theory of locally ordered higher dimensional structures), (2) simplicial sets (to obtain a theory of higher dimensional structures which come together with a simplicial enrichment at the highest level), (3) the category of sheaves on a locally connected space, or more generally a locally connected Grothendieck topos, (4) the algebras of any n -operad or (5) the category of multicategories (symmetric or not). The point is, the theory as we have developed it is actually *simpler* than before, and the generalisations mentioned here come at *no* extra cost.

Notation and terminology. Given a monad T on a category V the forgetful functor from the category of Eilenberg-Moore algebras of T is denoted as $U^T : V^T \rightarrow V$. We denote a T -algebra as a pair (X, x) , where X is the underlying object and $x : TX \rightarrow X$ is the algebra structure. For the category of globular sets it is typical to denote the image of the yoneda embedding as

$$0 \rightrightarrows 1 \rightrightarrows 2 \rightrightarrows 3 \rightrightarrows \dots$$

¹In this work we use the notion of contractibility given in [21] rather than the original notion of [3].

but then 0 denotes the globular set with one vertex and no edges or higher cells. Thus we adopt the convention of using 0 to denote objects of categories that we wish to think of as representing some underlying objects functor. Since initial objects are also important for us, we use the notation $\emptyset$ to denote them. While multicategories aren't directly multitensors, they become so after convolution – see [14]. Moreover when working seriously with multitensors, one is always manipulating functors of many variables, and so in fact working inside the **CAT**-enriched multicategory of categories. It is for these reasons that we find the term “multitensor” appropriate.

2. Connected objects and locally c-presentable categories

2.1. Connected objects in lextensive categories. We now collect together the basic, mostly well-known, abstract categorical theory of connected objects and coproduct decompositions. Recall that an object C in a category V with coproducts is *connected* when the representable $V(C, -)$ preserves coproducts.

The natural environment within which to study coproduct decompositions is a lextensive category. Recall that a category V is *extensive* when it has coproducts and for all families $(X_i : i \in I)$ of objects of V , the functor

$$\coprod : \prod_i (V/X_i) \rightarrow V / (\coprod_i X_i) \quad (f_i : Y_i \rightarrow X_i) \mapsto \coprod_i f_i : \coprod_i Y_i \rightarrow \coprod_i X_i$$

is an equivalence of categories. Note that this terminology is not quite standard: extensivity is usually defined using only finite coproducts.

Recall that coproducts in a category are said to be *disjoint* when coproduct coprojections are mono and the pullback of different coprojections is initial. Recall also that an initial object is said to be *strict* when any map into it is an isomorphism. The fundamental result on extensive categories is

Theorem 1. ([9],[12]) *A category V is extensive iff it has coproducts, pullbacks along coproduct coprojections and given a family of commutative squares*

$$\begin{array}{ccc} X_i & \xrightarrow{c_i} & X \\ f_i \downarrow & & \downarrow f \\ Y_i & \xrightarrow{d_i} & Y \end{array}$$

where $i \in I$ such that the d_i form a coproduct cocone, the c_i form a coproduct cocone iff these squares are all pullbacks. In an extensive category coproducts are disjoint and the initial object of V is strict.

A sufficient condition for extensivity is provided by

Lemma 2. *If a category V has coproducts, pullbacks along coproduct coprojections, and every $X \in V$ is a coproduct of connected objects, then V is extensive.*

PROOF. Suppose $f : X \rightarrow \emptyset$ is a morphism into the initial object, and $X = \coprod_{i \in I} X_i$ is a decomposition of X as a coproduct of connected objects. Then for any $i \in I$, one has by composing with c_i the i -th coproduct coprojection, a map $X_i \rightarrow \emptyset$. But since X_i is connected there can be no such map since the hom $V(X_i, \emptyset)$ is empty, and so I must be empty, and so X is initial, and so f is invertible. Thus V has a strict initial object.

Given A and B in V , denote by $c_A : A \rightarrow A + B$ the coprojection. Given a pair of maps $f, g : X \rightarrow A$ such that $c_A f = c_A g$, using X 's coproduct decomposition

again one has $c_A f c_i = c_A g c_i$, but since X_i is connected $f c_i = g c_i$, and since this is true for all i , $f = g$, and so c_A is mono. On the other hand suppose that a commutative square

$$\begin{array}{ccc} X & \longrightarrow & B \\ \downarrow & & \downarrow c_B \\ A & \xrightarrow{c_A} & A + B \end{array}$$

is given. Then by composing with c_i , one obtains another with X_i in place of X , but this cannot be since X_i is connected. Thus I is empty, and so X is initial. Thus the coproducts in V are disjoint.

By theorem(1) it suffices to show that given a family of commutative squares as on the left in

$$\begin{array}{ccc} X_i & \xrightarrow{c_i} & X \\ f_i \downarrow & & \downarrow f \\ Y_i & \xrightarrow{d_i} & Y \end{array} \qquad \begin{array}{ccc} V(Z, X_i) & \xrightarrow{V(Z, c_i)} & V(Z, X) \\ V(Z, f_i) \downarrow & & \downarrow V(Z, f) \\ V(Z, Y_i) & \xrightarrow{V(Z, d_i)} & V(Z, Y) \end{array}$$

where $i \in I$ such that the d_i form a coproduct cocone, the c_i form a coproduct cocone iff these squares are all pullbacks. By the yoneda lemma, this is equivalent to the same statement for the family of squares on the right (in **Set**) for all $Z \in V$. Since every object of V is a coproduct of connected ones, it suffices to consider just those $Z \in V$ that are connected. By the definition of connectedness, the functions $V(Z, c_i)$ (resp. $V(Z, d_i)$) form a coproduct cocone for all connected Z iff the maps c_i (resp d_i) do so in V , and so the result follows by the extensivity of **Set**. $\square$

Note that the conditions of lemma(2) are not necessary: there are many extensive categories whose objects don't decompose into coproducts of connected objects, for example, take the topos of sheaves on a space which is not locally connected. A category is *lexextensive* when it is extensive and has finite limits. There are many examples of lexextensive categories: Grothendieck toposes, the category of algebras of any higher operad and the category of topological spaces and continuous maps are all lexextensive.

We shall now study the decomposability of objects in such categories. As we shall see, the categorical datum which tells us whether all objects in a lexextensive category V admit a coproduct decomposition, is the left adjoint $(-) \cdot 1$ to the representable $V(1, -)$, where 1 as usual denotes the terminal object.

Lemma 3. *If V is lexextensive, then the representable $V(1, -) : V \rightarrow \mathbf{Set}$ has a left exact left adjoint $(-) \cdot 1$ given by taking copowers with 1 .*

PROOF. It is a standard fact, coming from nothing more than the universal property of coproducts, that the left adjoint take this form, and clearly $(-) \cdot 1$ preserves the terminal object. Given a pullback in **Set** as on the left in

$$\begin{array}{ccc} P & \xrightarrow{f} & B \\ h \downarrow & pb & \downarrow k \\ A & \xrightarrow{g} & C \end{array} \qquad \begin{array}{ccccc} h^{-1}(a) \cdot 1 & \longrightarrow & P \cdot 1 & \xrightarrow{f \cdot 1} & B \cdot 1 \\ \downarrow & pb & \downarrow h \cdot 1 & & \downarrow k \cdot 1 \\ 1 & \xrightarrow{a} & A \cdot 1 & \xrightarrow{g \cdot 1} & C \cdot 1 \end{array}$$

one has for each $a \in A$ a diagram as on the right. Since the original square is a pullback one has canonical bijections $h^{-1}(a) \cong k^{-1}(ga)$ enabling one to identify the top horizontal composite as $(-) \cdot 1$ applied to the inclusion of the fibre $k^{-1}(ga)$, and so for all a these maps exhibit $B \cdot 1$ as a coproduct. By theorem(1) it follows that the composite square on the right is a pullback. Since this is true for all $a \in A$ the right-most square is a pullback, again by theorem(1), as required. $\square$

This is very familiar in the case where V is a Grothendieck topos. Then the adjoint pair $(-) \cdot 1 \dashv V(1, -)$ is the global sections geometric morphism. We now note that the connectedness of an object in a lextensive category can be characterised in various ways.

Lemma 4. *Let V be a lextensive category and C be an object therein. Then C is connected iff the representable $V(C, -)$ preserves all copowers with 1. If in addition $(-) \cdot 1$ admits a left adjoint π_0 , then these conditions are equivalent to saying that $\pi_0(C) \cong 1$.*

PROOF. Suppose that $V(C, -)$ preserves copowers with 1. Coproduct coprojections defining $X = \coprod_{i \in I} X_i$ assemble, by theorem(1), into pullback squares

$$\begin{array}{ccc} X_i & \xrightarrow{c_i} & X \\ \downarrow & \text{pb} & \downarrow \\ 1 & \xrightarrow{i \cdot 1} & I \cdot 1 \end{array}$$

to which we apply $V(C, -)$. By theorem(1) in the case $V = \mathbf{Set}$, the functions $V(C, c_i)$ form a coproduct cocone since $V(C, i)$ do by hypothesis. Thus $V(C, -)$ does indeed preserve all coproducts. In the case where one has π_0 , by the canonical isomorphisms $V(C, I \cdot 1) \cong \mathbf{Set}(\pi_0 C, I)$, the connectedness of C is equivalent to $\mathbf{Set}(\pi_0 C, -)$ being isomorphic to the identity, which by the yoneda lemma is equivalent to $\pi_0 C \cong 1$. $\square$

We now characterise those lextensive categories in which every object admits as decomposition as a coproduct of connected objects.

Proposition 5. *Let V be a lextensive category. Then every object of V can be expressed as a coproduct of connected objects iff the functor $(-) \cdot 1$ admits a left adjoint, which we denote as π_0 . Under this hypothesis, a coproduct decomposition of a given object is unique up to canonical isomorphism.*

PROOF. Suppose that every object of V can be expressed as a coproduct of connected objects. For each $X \in V$ choose such a decomposition, write $\pi_0(X)$ for the indexing set, and for $i \in \pi_0(X)$ denote by $c_i : X_i \rightarrow X$ the corresponding coprojection. One induces the map η_X as in

$$(1) \quad \begin{array}{ccccc} X_i & \xrightarrow{c_i} & X & \searrow f & \\ \downarrow & \text{pb} & \downarrow \eta_X & & \\ 1 & \xrightarrow{i \cdot 1} & \pi_0(X) \cdot 1 & \xrightarrow{g \cdot 1} & I \cdot 1 \\ & & \searrow g(i) \cdot 1 & & \end{array}$$

so that the square commutes, and as indicated this square is a pullback by theorem(1). Given a set I and a morphism f as in (1), the connectedness of X_i ensures that there is a unique $g(i) \in I$ making the outside of (1) commute. In this way we have exhibited a unique $g : \pi_0 X \rightarrow I$ making the triangle in (1) commutative, and this exhibits η_X as the component at X of a unit of $\pi_0 \dashv (-) \cdot 1$. Moreover the uniqueness of coproduct decompositions is now evident, since any choice of all them gives rise in this way to an explicit left adjoint π_0 of the same functor $(-) \cdot 1$, and so different choices give rise to canonical isomorphisms between the corresponding π_0 's, which are compatible with the corresponding units.

For the converse let us suppose that we have $\pi_0 \dashv (-) \cdot 1$. Then one has $\eta_X : X \rightarrow \pi_0(X) \cdot 1$, and one then takes pullbacks as in (1) to obtain the $c_i : X_i \rightarrow X$ which form a coproduct cocone by theorem(1). To finish the proof we must show that all these X_i 's are connected, and by lemma(4) it suffices to show that the cardinality $|\pi_0(X_i)|$ is 1. If it was 0 then one would have $\eta_{X_i} : X_i \rightarrow \emptyset$ making X_i initial too, since initial objects are strict. But then by defining $I = \pi_0(X) \setminus \{i\}$, writing $j : I \rightarrow \pi_0(X)$ for the proper inclusion, one has also $\eta' : X \rightarrow I \cdot 1$ such that $(j \cdot 1)\eta' = \eta_X$. But by the universal property of η_X one also has a section $s : \pi_0(X) \rightarrow I$ of j , contradicting the properness of j . Thus $|\pi_0(X_i)| > 0$. Note that we have a diagram

$$\begin{array}{ccccc}
 & & X_i & \xrightarrow{c_i} & X \\
 & \nearrow \eta_{X_i} & \downarrow & & \downarrow \eta_X \\
 & & 1 & \xrightarrow{i} & \pi_0(X) \cdot 1 \\
 \pi_0(X_i) \cdot 1 & \nearrow & \text{(I)} & \searrow & \\
 & & \pi_0(c_i) \cdot 1 & &
 \end{array}$$

in which the outside and all regions except region (I) are clearly commutative. By the universal property of η_{X_i} the function $\pi_0(c_i)$ factors as

$$\pi_0(X_i) \longrightarrow 1 \xrightarrow{i} \pi_0(X)$$

so that $|\text{im}(\pi_0(c_i))| \leq 1$, and since $\pi_0(c_i)$ as a coprojection in **Set** is injective, we have $|\pi_0(X_i)| \leq 1$. $\square$

2.2. Locally c-presentable categories. We recall first some of the basic notions from the theory of locally presentable categories [2, 15, 22]. Let λ be a regular cardinal. A λ -small category is one whose class of arrows forms a set of cardinality $< \lambda$, and a category $\mathcal{A}$ is λ -filtered when every functor $J \rightarrow \mathcal{A}$, where J is λ -small, admits a cocone. Colimits of functors out of λ -filtered categories are called λ -filtered colimits. An object X of a category V is λ -presentable when the representable $V(X, -)$ preserves all λ -filtered colimits that exist in V . A locally small category V is *locally λ -presentable* when it is cocomplete and there is a set $\mathcal{S}$ of λ -presentable objects such that every object of V is a λ -filtered colimit of objects from $\mathcal{S}$.

There are many alternative characterisations of locally λ -presentable categories, the most minimalistic being the following. Recall that a set $\mathcal{D}$ of objects of V is a *strong generator* when for all maps $f : X \rightarrow Y$, if

$$V(\mathcal{D}, f) : V(\mathcal{D}, X) \rightarrow V(\mathcal{D}, Y)$$

is bijective for all $D \in \mathcal{D}$, then f is an isomorphism. Then a locally small category V is locally λ -presentable iff it is cocomplete and has a strong generator consisting of λ -presentable objects. Other characterisations include: as categories of **Set**-valued models of limit sketches whose distinguished cones are λ -small; as full reflective subcategories of presheaf categories for which the inclusion is λ -accessible; to name just two. See for instance [2, 15, 22] for a complete discussion of this fundamental notion.

The appropriate functors between such categories are the λ -accessible ones – a functor being λ -accessible when it preserves λ -filtered colimits. Here we describe a mild variant of these notions in which the role of λ -presentable objects is played by objects which are both λ -presentable and connected, and accessible functors are replaced by functors which preserve both λ -filtered colimits and coproducts. A category $\mathcal{A}$ is λ -c-filtered when every functor $J \rightarrow \mathcal{A}$, where J is λ -small and connected, admits a cocone. Clearly a category is λ -c-filtered iff its connected components are λ -filtered, and thus a λ -c-filtered colimit is the same thing as a coproduct of λ -filtered colimits. The categories which are λ -small and connected form a *doctrine* $\mathbb{D}$ in the sense of [1], a λ -c-filtered category is one which is $\mathbb{D}$ -filtered in the sense of [1] definition(1.2), and this doctrine is easily exhibited as *sound* in the sense of [1] definition(2.2).

Definition 6. [1] A locally small category V is *locally λ -c-presentable* when it is cocomplete and has a set $\mathcal{S}$ of objects which are connected and λ -presentable, such that every object of V is a λ -c-filtered colimit of objects of $\mathcal{S}$. A *locally c-presentable* category is one which is locally λ -c-presentable for some regular cardinal λ . When λ is the first infinite ordinal, we also use the terminology locally *finitely* c-presentable.

We have attributed definition(6) to [1] since a locally λ -c-presentable category is exactly a locally- $\mathbb{D}$ -presentable category in the sense of [1] definition(5.1), for the doctrine $\mathbb{D}$ of λ -small connected categories. In theorem(8) below we collect together the various reformulations of this notion.

Lemma 7. *Let λ be a regular cardinal and V be locally λ -presentable and extensive. Then a summand of a λ -presentable object in V is λ -presentable.*

PROOF. Suppose that $A, B \in V$ and that their coproduct $A + B$ is a λ -presentable object. Since V is locally λ -presentable one has a λ -filtered category I , and a colimit cocone $k_i : A_i \rightarrow A$ for $i \in I$, where the A_i are λ -presentable objects. Thus the maps $k_i + 1_B : A_i + B \rightarrow A + B$ exhibit $A + B$ as a λ -filtered colimit. Since $A + B$ is λ -presentable, there is $j \in I$ and $s : A + B \rightarrow A_j + B$ such that $(k_j + 1_B)s = 1_{A+B}$. Extensivity ensures that the right-most square in

$$\begin{array}{ccccc}
 & & 1 & & \\
 & \curvearrowright & & \curvearrowright & \\
 A & \xrightarrow{\quad t \quad} & A_j & \xrightarrow{\quad k_j \quad} & A \\
 \downarrow c_A & & \downarrow c_{A_j} & & \downarrow c_A \\
 A + B & \xrightarrow{\quad s \quad} & A_j + B & \xrightarrow{\quad k_j + 1_B \quad} & A + B \\
 & \curvearrowleft & & \curvearrowleft & \\
 & & 1 & &
 \end{array}$$

is a pullback, enabling us to induce t as shown which exhibits k_j as a retraction, and thus A as λ -presentable. $\square$

Theorem 8. *For a locally small category V and regular cardinal λ , the following statements are equivalent.*

- (1) V is cocomplete and has a strong generator consisting of objects which are connected and λ -presentable.
- (2) V is locally λ -c-presentable.
- (3) V is cocomplete and has a small dense subcategory consisting of objects which are connected and λ -presentable.
- (4) V is a full subcategory of a presheaf category for which the inclusion is λ -accessible, coproduct preserving and has a left adjoint.
- (5) V is equivalent to the category of models for a limit sketch whose distinguished cones are λ -small and connected.
- (6) V is equivalent to the full subcategory of $[\mathcal{A}, \mathbf{Set}]$ consisting of λ -small connected limit preserving functors, for some small category $\mathcal{A}$ with λ -small connected limits.
- (7) V is a free completion of a small category with λ -small connected limits under λ -c-filtered colimits.
- (8) V is locally λ -presentable and every object of V is a coproduct of connected objects.
- (9) V is locally λ -presentable, extensive and the functor $(-)\cdot 1 : \mathbf{Set} \rightarrow V$ has a left adjoint.

PROOF. The equivalence of (2), (5), (6) and (7) is [1] theorem(5.5) in the case where $\mathbb{D}$ is the doctrine for λ -small connected categories. The implication (3) $\Rightarrow$ (1) is trivial, and the equivalence of (8) and (9) is an immediate consequence of lemma(2) and proposition(5). Given (2) V is clearly locally λ -presentable and any $X \in V$ is a coproduct of λ -filtered colimits of λ -presentable connected objects. But a λ -filtered colimit of connected objects is connected, and so (2) $\Rightarrow$ (8).

(1) $\Rightarrow$ (2): Let $\mathcal{D}$ be a strong generator of λ -small connected objects, and denote also by $\mathcal{D}$ the full subcategory of V it determines. Take the closure $\mathcal{S}$ of $\mathcal{D}$ in V under λ -small connected colimits, and note that $\mathcal{S}$ is also essentially small (see [17] section(3.5)). Thus $\mathcal{S}$ is also a strong generator of V consisting of λ -small connected objects and moreover, the full subcategory it determines has λ -small connected colimits. Thus for $X \in V$, the comma category $\mathcal{S}/X$ is λ -c-filtered, and so it suffices to show that the comma object defining $\mathcal{S}/X$ exhibits X as a colimit. Denote by K the actual colimit, for $f : A \rightarrow X$ in $\mathcal{S}/X$ by $\kappa_f : A \rightarrow K$ the component of the colimit cocone, and by $k : K \rightarrow X$ the induced map. Since $\mathcal{S}$ is a strong generator it suffices to show that for all $A \in \mathcal{S}$ the function $V(A, k) : V(A, C) \rightarrow V(A, X)$ is bijective. It is surjective by the definition of k , which is defined as the unique map such that $k\kappa_f = f$ for all $f \in \mathcal{S}/X$. To see that $V(A, k)$ is injective, suppose that one has b and $c : A \rightarrow K$ such that $kb = kc$. Then since the colimit defining K is λ -c-filtered and A is λ -presentable and connected, one has $b_2 : B \rightarrow K$ and $b_3 : B \rightarrow X$ such that $\kappa_{b_3}b_2 = b$, and similarly $c_2 : C \rightarrow K$ and $c_3 : C \rightarrow X$ such that $\kappa_{c_3}c_2 = c$. Take the pushout

$$\begin{array}{ccc} A & \xrightarrow{b_2} & B \\ c_2 \downarrow & po & \downarrow p \\ C & \xrightarrow{q} & D \end{array}$$

in $\mathcal{S}$, and induce $d : D \rightarrow X$ as the unique map such that $dp = b_3$ and $dq = c_3$. The result follows by

$$b = \kappa_{b_3} b_2 = \kappa_d p b_2 = \kappa_d q c_2 = \kappa_{c_3} c_2 = c.$$

(2) $\Rightarrow$ (3): Let $\mathcal{S}$ be the set of λ -presentable connected objects required by definition(6), and denote by $i : \mathcal{S} \rightarrow V$ the inclusion of the corresponding full subcategory of V . Let $X, Y \in V$ and suppose that $\phi : V(i, X) \rightarrow V(i, Y)$ in $\widehat{\mathcal{S}}$ is given. One has $k : J \rightarrow \mathcal{S}$ with J small and λ -c-filtered, such that $\text{col}(ik) = X$, and we denote by $\kappa_j : kj \rightarrow X$ a typical component of the colimiting cocone. Induce $\phi' : X \rightarrow Y$ as the unique map such that $\phi' \kappa_j = \phi(\kappa_j)$ for all $j \in J$. But then for all $f : S \rightarrow X$ with $S \in \mathcal{S}$, one has $\phi' f = \phi(f)$: since S is λ -presentable and connected one can find $j \in J$ and $g : S \rightarrow kj$ such that $f = \kappa_j g$ and so

$$\phi(f) = \phi(\kappa_j)g = \phi' \kappa_j g = \phi' f.$$

Thus $V(i, 1) : V \rightarrow \widehat{\mathcal{S}}$ is fully-faithful, in other words, i is dense as claimed.

(3) $\Rightarrow$ (4): Let $i : \mathcal{S} \rightarrow V$ be the inclusion of the given dense subcategory. Then $V(i, 1) : V \rightarrow \widehat{\mathcal{S}}$ preserves coproducts and is λ -accessible since the objects of $\mathcal{S}$ are connected and λ -presentable, is fully faithful since i is dense, and has a left adjoint given by left kan extension along i since V is cocomplete and $\mathcal{S}$ is small.

(4) $\Rightarrow$ (3): Let $I : V \rightarrow \widehat{\mathcal{C}}$ be the given inclusion and L be its left adjoint. Let T be the monad induced by $L \dashv I$, and note that since I is fully-faithful, it is monadic. Denote by $i : \mathcal{S} \rightarrow \widehat{\mathcal{C}}$ the inclusion of the closure of the representables in $\widehat{\mathcal{C}}$ under λ -small connected colimits. Since the objects of $\mathcal{S}$ are λ -presentable and connected, and T is λ -accessible and coproduct preserving, it follows that $(T, \mathcal{S})$ is a monad with arities in the sense of [26, 23, 8]. Taking

$$\mathcal{S} \xrightarrow{k} \Theta_T \xrightarrow{j} V$$

to be the identity on objects fully faithful factorisation of Li , it follows from the Nerve theorem [8] that $j : \Theta_T \rightarrow V$ is dense. Since for all $A \in \Theta_T$, $V(jA, -) \cong V(LiA, -) \cong \widehat{\mathcal{C}}(iA, IA)$ and I preserves coproducts and λ -filtered colimits, it follows that the image of j consists of connected λ -presentable objects.

(9) $\Rightarrow$ (1): V is cocomplete by definition. Let $\mathcal{D}$ be a strong generator of λ -presentable objects of V . Decompose each object of $\mathcal{D}$ into connected components using proposition(5), and write $\mathcal{D}'$ for the set of summands of objects of $\mathcal{D}$ that so arise. Clearly $\mathcal{D}'$ is also a strong generator of V , its objects are connected by definition and λ -presentable by lemma(7). $\square$

Examples 9. By theorem(8)(4) any presheaf topos is locally finitely c-presentable. By theorem(8)(9) a Grothendieck topos is locally connected iff it is locally c-presentable.

Just as with locally presentable categories, locally c-presentable categories are closed under many basic categorical constructions. For instance from theorem(8)(8), one sees immediately that the slices of a locally λ -c-presentable category are locally λ -c-presentable from the corresponding result for locally presentable categories. Another instance of this principle is the following result.

Theorem 10. *If V is locally λ -c-presentable and T is a λ -accessible coproduct preserving monad on V , then V^T is locally λ -c-presentable.*

PROOF. By the analogous result for locally presentable categories V^T is locally λ -presentable and thus cocomplete. Defining Θ_0 to be the full subcategory of V consisting of the λ -presentable and connected objects, (T, Θ_0) is a monad with arities in the sense of [26]. One has a canonical isomorphism

$$\begin{array}{ccc} V^T & \xrightarrow{V^T(i,1)} & \hat{\Theta}_T \\ U^T \downarrow & \cong & \downarrow \text{res}_j \\ V & \xrightarrow{V(i_0,1)} & \hat{\Theta}_0 \end{array}$$

in the notation of [26]. Since T preserves λ -filtered colimits and coproducts, U^T creates them. Since j is bijective on objects res_j creates all colimits. Thus by the above isomorphism $V^T(i, 1)$ preserves λ -filtered colimits and coproducts. By the nerve theorem of [26] $V^T(i, 1)$ is also fully faithful, it has a left adjoint since V^T is cocomplete given by left extending i along the yoneda embedding, and so we have exhibited V^T as conforming to theorem(8)(4). $\square$

Examples 11. An n -operad for $0 \leq n \leq \omega$ in the sense of [3], gives a finitary coproduct preserving monad on the category $\hat{\mathbb{G}}_{\leq n}$ of n -globular sets, and its algebras are just the algebras of the monad. Since $\hat{\mathbb{G}}_{\leq n}$ as a presheaf topos is locally finitely c-presentable by example(9), the category of algebras of any n -operad is locally finitely c-presentable by theorem(10).

3. Categories of enriched graphs

3.1. Enriched graphs.

Definition 12. Let V be a category. A *graph X enriched in V* consists of an underlying set X_0 whose elements are called *objects*, together with an object $X(a, b)$ of V for each ordered pair (a, b) of objects of X . The object $X(a, b)$ will sometimes be called the *hom* from a to b . A morphism $f : X \rightarrow Y$ of V -enriched graphs consists of a function $f_0 : X_0 \rightarrow Y_0$ together with a morphism $f_{a,b} : X(a, b) \rightarrow Y(fa, fb)$ for each (a, b) . The category of V -graphs and their morphisms is denoted as $\mathcal{GV}$, and we denote by $\mathcal{G}$ the obvious 2-functor

$$\mathcal{G} : \mathbf{CAT} \rightarrow \mathbf{CAT} \quad V \mapsto \mathcal{GV}$$

with object map as indicated.


Note that for $n \in \mathbb{N}$, $\mathcal{G}^n \mathbf{Set}$ is the category of n -globular sets, and that $\mathcal{G} \mathbf{Glob} \cong \mathbf{Glob}$ where $\mathbf{Glob}$ denotes the category of globular sets. In fact applying the 2-functor $\mathcal{G}$ successively to the inclusion of the empty category into the point (ie the terminal category), one obtains the inclusion of the category of $(n-1)$ -globular sets into the category of n -globular sets. In the case $n > 0$ this is the inclusion with object map

$$X_0 \xleftarrow{\quad} \cdots \xleftarrow{\quad} X_{n-1} \quad \mapsto \quad X_0 \xleftarrow{\quad} \cdots \xleftarrow{\quad} X_{n-1} \xleftarrow{\quad} \emptyset$$

and when $n=0$ this is the functor $1 \rightarrow \mathbf{Set}$ which picks out the empty set. Thus there is exactly one (-1) -globular set which may be identified with the empty set.

When V has an initial object $\emptyset$, one can regard any sequence of objects $(Z_1, \dots, Z_n)$ of V as a V -graph. The object set is $\{0, \dots, n\}$, $(Z_1, \dots, Z_n)(i-1, i) = Z_i$ for $1 \leq i \leq n$,

and all the other homs are equal to $\emptyset$. We denote also by 0 the V -graph corresponding to the empty sequence $()$. Note that 0 is a representing object for the forgetful functor $(-)_0 : \mathcal{G}V \rightarrow \mathbf{Set}$ which sends an enriched graph to its underlying set of objects. Globular pasting diagrams [3] may be regarded as iterated sequences, for instance $(0, 0, 0)$ and $((0, 0), (0), (0, 0, 0))$ correspond respectively to



when one starts with $V = \mathbf{Set}$. We denote by “ n ” the free-living n -cell, defined inductively by $n + 1 = (n)$.

It is often better to think of $\mathcal{G}$ as taking values in $\mathbf{CAT}/\mathbf{Set}$. By applying the endofunctor $\mathcal{G}$ to the unique functor $t_V : V \rightarrow 1$ for each V , produces $(-)_0$ which sends an enriched graph to its underlying set of objects. This 2-functor

$$\mathcal{G}_1 : \mathbf{CAT} \rightarrow \mathbf{CAT}/\mathbf{Set}$$

has a left adjoint described as follows. First to a given functor $f : A \rightarrow \mathbf{Set}$ we denote by $f^{\times 2} : A \rightarrow \mathbf{Set}$ the functor with object map $a \mapsto f(a) \times f(a)$. Then to a given functor $g : A \rightarrow \mathbf{Set}$ we denote by $g_\bullet$ the domain of the discrete opfibration corresponding to g via the Grothendieck construction. That is, $g_\bullet$ can be defined via the pullback

$$\begin{array}{ccc} g_\bullet & \longrightarrow & \mathbf{Set}_\bullet \\ \downarrow & \text{\scriptsize } pb & \downarrow U \\ A & \xrightarrow{g} & \mathbf{Set} \end{array}$$

where U is the forgetful functor from the category of pointed sets and base point preserving maps. The left adjoint to $\mathcal{G}_1$ is then described on objects by $f \mapsto f_\bullet^{\times 2}$. Explicitly $f_\bullet^{\times 2}$ has as objects triples (a, x, y) where a is an object of A , and (x, y) is an ordered pair of objects of $f a$. Maps in $f_\bullet^{\times 2}$ are maps in A which preserve these base points.

It is interesting to look at the unit and counit of this 2-adjunction. Given a category V , $(\mathcal{G}t_V)_\bullet^{\times 2}$ is the category of bipointed enriched graphs in V . The counit $\varepsilon_V : (\mathcal{G}t_V)_\bullet^{\times 2} \rightarrow V$ sends (X, a, b) to the hom $X(a, b)$. When V has an initial object ε_V has a left adjoint L_V given by $X \mapsto ((X), 0, 1)$. Given a functor $f : A \rightarrow \mathbf{Set}$ the unit $\eta_f : A \rightarrow \mathcal{G}(f_\bullet^{\times 2})$ sends $a \in A$ to the enriched graph whose objects are elements of $f a$, and the hom $\eta_f(a)(x, y)$ is given by the bipointed object (a, x, y) .

A given functor $f : A \rightarrow \mathcal{G}V$ thus admits a canonical factorisation

$$A \xrightarrow{\eta_{f_0}} \mathcal{G}(f_{0\bullet}^{\times 2}) \xrightarrow{\mathcal{G}\mathcal{H}_V(f)} \mathcal{G}V$$

where on objects one has $\mathcal{H}_V(a, x, y) = f(a)(x, y)$. This is the generic factorisation of f in the sense of [26]. The adjointness $(-)_\bullet^{\times 2} \dashv \mathcal{G}_1$ says that f is uniquely determined by its object part $f_0 := (-)_0 f$ and its hom data $\mathcal{H}_V(f)$. For the sake of computing colimits in $\mathcal{G}V$, as we will in section(3.2), it is worth noting that one

can reorganise the data of a lax triangle as on the left in

$$\begin{array}{ccc}
 \begin{array}{ccc} A & \xrightarrow{k} & B \\ & \searrow f & \swarrow h \\ & \phi & \\ & \mathcal{G}V & \end{array} &
 \begin{array}{ccc} A & \xrightarrow{k} & B \\ & \searrow f_0 & \swarrow h_0 \\ & \phi_0 & \\ & \mathbf{Set} & \end{array} &
 \begin{array}{ccc} f_{0\bullet}^{\times 2} & \xrightarrow{\phi_{0\bullet}^{\times 2}} & h_{0\bullet}^{\times 2} \\ & \searrow \mathcal{H}_V(f) & \swarrow \mathcal{H}_V(h) \\ & \mathcal{H}_\phi & \\ & V & \end{array}
 \end{array}$$

into $\mathcal{G}V$ in the same way. The middle triangle is just $(-)_0\phi$. In the right hand triangle, $\phi_{0\bullet}^{\times 2}$ is the evident functor with object map $(a, x, y) \mapsto (ka, \phi_a(x), \phi_a(y))$ which is determined by ϕ_0 . The natural transformation $\mathcal{H}_\phi$ has components given by the hom maps of the components of ϕ , that is $(\mathcal{H}_\phi)_{(a,x,y)}$ is the map $(\phi_a)_{x,y} : f(a)(x, y) \rightarrow hk(a)(\phi_a(x), \phi_a(y))$. It then follows easily from unpacking the data involved that

Lemma 13. *Given $f : A \rightarrow \mathcal{G}V$, $k : A \rightarrow B$ and $h : B \rightarrow \mathcal{G}V$, the assignment $\phi \mapsto (\phi_0, \mathcal{H}_\phi)$ is a bijection which is natural in h .*

Suppose that one has an object 0 in a category A , and f is the representable $f = A(0, -)$. Then $f_{\bullet}^{\times 2}$ may be regarded as the category of endo-cospans of the object 0, that is to say the category of diagrams

$$0 \rightarrow a \leftarrow 0$$

and a point of $a \in A$ is now just a map $0 \rightarrow a$. When A is also cocomplete one can compute a left adjoint to η_f . To do this note that a graph X enriched in $f_{\bullet}^{\times 2}$ gives rise to a functor

$$\overline{X} : X_0^{(2)} \rightarrow A$$

where X_0 is the set of objects of X . For any set Z , $Z^{(2)}$ is defined as the following category. It has two kinds of objects: an object being either an element of Z , or an ordered pair of elements of Z . There are two kinds of non-identity maps

$$x \rightarrow (x, y) \leftarrow y$$

where (x, y) is an ordered pair from Z , and $Z^{(2)}$ is free on the graph just described. A more conceptual way to see this category is as the category of elements of the graph

$$Z \times Z \rightrightarrows Z$$

where the source and target maps are the product projections, as a presheaf on the category

$$\mathbb{G}_{\leq 1} = \boxed{0 \rightrightarrows 1}$$

and so there is a discrete fibration $Z^{(2)} \rightarrow \mathbb{G}_{\leq 1}$. The functor $\overline{X}$ sends singletons to $0 \in A$, and a pair (x, y) to the head of the hom $X(x, y)$. The arrow map of $\overline{X}$ encodes the bipointings of the homs. One may then easily verify

Proposition 14. *Let $0 \in A$, $f = A(0, -)$ and A be cocomplete. Then η_f has left adjoint given on objects by $X \mapsto \text{colim}(\overline{X})$.*

There is a close connection between $\mathcal{G}$ and the Fam construction. A very mild reformulation of the notion of V -graph is the following: a V -graph X consists of a set X_0 together with an $(X_0 \times X_0)$ -indexed family of objects of V . Together with

the analogous reformulation of the maps of $\mathcal{G}V$, this means that we have a pullback square

$$\begin{array}{ccc} \mathcal{G}V & \longrightarrow & \mathbf{Fam}V \\ (-)_0 = \mathcal{G}t_V \downarrow & \lrcorner & \downarrow \mathbf{Fam}(t_V) \\ \mathbf{Set} & \xrightarrow{(-)^2} & \mathbf{Set} \end{array}$$

in **CAT**, and thus a cartesian 2-natural transformation $\mathcal{G} \rightarrow \mathbf{Fam}$. From [26] theorem(7.4) we conclude

Proposition 15. *$\mathcal{G}$ is a familial 2-functor.*

In particular notice that for all V , the functor $(-)_0 : \mathcal{G}V \rightarrow \mathbf{Set}$ has the structure of a split fibration. The cartesian morphisms are exactly those which are *fully faithful*, which are those morphisms of V -graphs whose hom maps are isomorphisms. The vertical-cartesian factorisation of a given $f : X \rightarrow Y$ corresponds to its factorisation as an identity on objects map followed by a fully-faithful map. Moreover it follows from the theory of [26] that $\mathcal{G}$ preserves conical connected limits as well as all the notions of “Grothendieck fibration” which one can define internal to a finitely complete 2-category, and that the obstruction maps for comma objects are right adjoints. In addition to this we have

Lemma 16. *$\mathcal{G}$ preserves Eilenberg-Moore objects.*

Given a monad T on a category V , we shall write V^T for the category of T -algebras and morphisms thereof, and $U^T : V^T \rightarrow V$ for the forgetful functor. We shall denote a typical object of V^T as a pair (X, x) , where X is the underlying object in V and $x : TX \rightarrow X$ is the T -algebra structure. From [24] the 2-cell $TU^T \rightarrow U^T$, whose component at (X, x) is x itself has a universal property exhibiting V^T as a kind of 2-categorical limit called an *Eilenberg-Moore object*. See [24] or [20] for more details on this general notion. The direct proof that for any monad T on a category V , the obstruction map $\mathcal{G}(V^T) \rightarrow \mathcal{G}(V)^{\mathcal{G}(T)}$ is an isomorphism comes down to the obvious fact that for any V -graph B , a $\mathcal{G}T$ -algebra structure on B is the same thing as a T -algebra structure on the homs of B , and similarly for algebra morphisms.

Returning to the consideration of $\mathcal{G}_1$, our final observation for this section is

Proposition 17. *$\mathcal{G}_1 : \mathbf{CAT} \rightarrow \mathbf{CAT}/\mathbf{Set}$ is locally fully faithful.*

PROOF. Given functors $F, G : V \rightarrow W$, the data of a natural transformation $\phi : \mathcal{G}F \rightarrow \mathcal{G}G$ over **Set** amounts to giving for each $X \in \mathcal{G}V$ and $a, b \in X_0$, maps $\phi_{X,a,b} : FX(a, b) \rightarrow GX(a, b)$, such that for $f : X \rightarrow Y$ one has the naturality condition for f between a and b :

$$\begin{array}{ccc} FX(a, b) & \xrightarrow{\phi_{X,a,b}} & GX(a, b) \\ Ff_{a,b} \downarrow & = & \downarrow Gf_{a,b} \\ FY(fa, fb) & \xrightarrow{\phi_{Y,fa,fb}} & GY(fa, fb) \end{array}$$

So we define $\phi' : F \rightarrow G$ by $\phi'_Z = \phi_{(Z),0,1}$. One has $c : (X(a, b)) \rightarrow X$ in $\mathcal{G}V$ with object map $(0, 1) \mapsto (a, b)$ and hom map $c_{0,1}$ the identity. The naturality

condition for c between 0 and 1 yields $\phi_{(X(a,b)),0,1} = \phi_{X,a,b}$ from which it follows that $\phi = \mathcal{G}\phi'$. Conversely $(\mathcal{G}\phi)'_Z = (\mathcal{G}\phi)_{(Z),0,1} = \phi_Z$ and so $\phi \mapsto \phi'$ is the inverse of

$$\mathbf{CAT}(V, W)(F, G) \rightarrow \mathbf{CAT}/\mathbf{Set}(\mathcal{G}V, \mathcal{G}W)(\mathcal{G}F, \mathcal{G}G) \quad \psi \mapsto \mathcal{G}\psi. \quad \square$$

3.2. Properties of $\mathcal{G}V$. This section contains a variety of results from which it is clear that as a category, $\mathcal{G}V$ is at least as good as V . To begin with, any limit that V possesses is also possessed by $\mathcal{G}V$.

Proposition 18. *Let I be a small category. If V admits limits of functors out of I , then so does $\mathcal{G}V$ and these are preserved by $(-)_0$.*

PROOF. Let $F : I \rightarrow \mathcal{G}V$ be a functor. We construct its limit L directly as follows. First we take the set L_0 to be the limit of the functor $F(-)_0$, writing $\lambda_{i,0} : L_0 \rightarrow F(i)_0$ for a typical component of the limit cone. Without loss of generality one can represent the elements of L_0 explicitly as matching families of elements of the $F(i)_0$. That is, any such element is a family

$$x := (x_i \in F(i)_0 : i \in I)$$

such that for all $f : i \rightarrow j$ in I , one has $Ff(x_i) = x_j$. Given an ordered pair (x, y) of such families, one has a functor

$$F_{x,y} : I \rightarrow V \quad i \mapsto Fi(x_i, y_i)$$

with indicated object map. One then defines the hom $L(x, y)$ to be the limit in V of $F_{x,y}$, and we write $\lambda_{i,x,y} : Fi(x_i, y_i) \rightarrow L(x, y)$ for the components of the limit cone. These provide the hom maps, and $\lambda_{i,0}$ the object functions, of morphisms $\lambda_i : L \rightarrow Fi$. It is easily verified that these exhibit L as a limit of F . $\square$

In particular from the explicit construction of limits just described, it is clear that $\mathcal{G}V$ possesses some pullbacks under no conditions on V .

Corollary 19. *For any category V , $\mathcal{G}V$ admits all pullbacks along fully faithful maps, and these are preserved by $(-)_0$. Moreover the pullback of a fully faithful map is itself fully faithful.*

PROOF. In this case the construction of proposition(18) goes through because the pullbacks in V that arise in the construction are all along an isomorphism, and such clearly exist in any V . The last statement follows from this explicit construction since isomorphisms in any V are pullback stable. $\square$

By lemma(13) one can compute the left kan extension of $F : I \rightarrow \mathcal{G}V$, along any functor $G : I \rightarrow J$ between small categories, in the following way. First compute the left extension $K_0 : J \rightarrow \mathbf{Set}$ of F_0 along G denoting the universal 2-cell as $\kappa_0 : F_0 \rightarrow K_0G$. Then given sufficient colimits in V , compute the left extension $\mathcal{H}_V(K) : K_0^{\times 2} \rightarrow V$ of $\mathcal{H}_V(F)$ along $\kappa_0^{\times 2}$, denoting the universal 2-cell as $\mathcal{H}_\kappa : \mathcal{H}_V(F) \rightarrow \mathcal{H}_V(K)$. Thus we have the object part K_0 and hom data $\mathcal{H}_V(K)$ of a functor $K : J \rightarrow \mathcal{G}V$. The natural transformation $\kappa : F \rightarrow KG$ corresponding to $(\kappa_0, \mathcal{H}_\kappa)$ by lemma(13) clearly exhibits K as the left extension of F along G , by a straight forward application of lemma(13) and the definition of “left kan extension”.

When $J = 1$ note that $K_0^{\times 2}$ is just the discrete category $K_0 \times K_0$, and so for $x, y \in K_0$ one may compute $\mathcal{H}_V(K)(x, y)$ as the colimit of $\mathcal{H}_V(F)$ restricted to the fibre of $\kappa_0^{\times 2}$ over (x, y) .

Proposition 20. (1) *For any category V , $\mathcal{G}V$ has a strict initial object.*

- (2) If V has an initial object, then $\mathcal{G}V$ has coproducts.
- (3) If λ is a regular cardinal and V has λ -filtered colimits, then so does $\mathcal{G}V$.
- (4) If V has all small colimits, then so does $\mathcal{G}V$.

In each case the colimits in $\mathcal{G}V$ under discussion are preserved by $(-)_0$.

PROOF. By the above uniform construction of colimits one has (4), and the preservation of any colimit by $(-)_0$ when it is constructed in this way. The empty V -enriched graph is clearly strictly initial in $\mathcal{G}V$ and so (1) follows.

(2): In the case where I is discrete, $\kappa_{0,\bullet}^{\times 2}$ is the inclusion

$$\coprod_{i \in I} F(i)_0 \times F(i)_0 \rightarrow \coprod_i F(i)_0 \times \coprod_i F(i)_0$$

(between discrete categories) which picks out pairs (x, y) which live in the same component. Thus the coproduct $X := \coprod_i X_i$ in $\mathcal{G}V$ is defined to have objects those of the disjoint union of the objects sets of the X_i , and homs given by $X(x, y) = X_i(x, y)$ when x and y are both in X_i , and $\emptyset$ otherwise.

(3): When I is λ -filtered we note that since λ -filtered colimits in **Set** commute with binary products, the cocone

$$\kappa_{0,i} \times \kappa_{0,i} : F(i)_0 \times F(i)_0 \rightarrow K_0 \times K_0$$

in **Set** is also a colimit cocone. Thus the functor $\kappa_{0,\bullet}^{\times 2} : F_{0,\bullet}^{\times 2} \rightarrow K_0 \times K_0$ has another interpretation. Since $F_{0,\bullet}^{\times 2}$ is the category of elements of the functor $i \mapsto F(i)_0 \times F(i)_0$, then by the above remark $K_0 \times K_0$ is the set of connected components of $F_{0,\bullet}^{\times 2}$ and $\kappa_{0,\bullet}^{\times 2}$ is the canonical projection. So the fibres of $\kappa_{0,\bullet}^{\times 2}$ are the connected components of $F_{0,\bullet}^{\times 2}$. Since the evident forgetful functor $F_{0,\bullet}^{\times 2} \rightarrow I$ is a discrete opfibration, the connected components of $F_{0,\bullet}^{\times 2}$ are themselves λ -filtered. Thus a fibre of $\kappa_{0,\bullet}^{\times 2}$ over a given (x, y) will itself be λ -filtered, and so λ -filtered colimits in V will suffice for the construction of the colimit in this case. $\square$

Remark 21. There is one very easy to understand class of limit/colimit of V -graphs. These are those for functors $F : J \rightarrow \mathcal{G}V$ where J is connected and $F_0 : J \rightarrow \mathbf{Set}$ is constant at some set X . For then the limit or colimit of F may also be taken to have object set X , and one computes the hom between a and $b \in X$ of the limit or colimit by taking the limit or colimit in V of the functor $J \rightarrow V$ with object map $j \mapsto F(j)(a, b)$.

Let $X \in \mathcal{G}V$ and suppose that V has an initial object $\emptyset$. We consider first the relation on X_0 defined as follows

$$\{(a, b) : V(X(a, b), \emptyset) = \emptyset\} \subseteq X_0 \times X_0$$

and say that a and b are *in the same component* of X when they are identified by the equivalence relation generated by the above relation. Denote by $\pi_0(X)$ the set of equivalence classes, which themselves are called connected components. For $i \in \pi_0(X_0)$ we denote by X_i the full sub- V -graph of X whose objects are those of X 's i -th component, and by $c_{X,i} : X_i \rightarrow X$ the evident inclusion. The assignment $X \mapsto \pi_0(X)$ is the object map of a functor $\pi_0 : \mathcal{G}V \rightarrow \mathbf{Set}$.

Proposition 22. *Let V have an initial object $\emptyset$ and $X \in \mathcal{G}V$. Then X is connected iff $\pi_0(X) = 1$.*

PROOF. Suppose that $\pi_0(X) = 1$. We must show that any map $f : X \rightarrow \coprod_i Y_i$ factors uniquely through a unique summand. From the explicit construction of coproducts in $\mathcal{G}V$ in proposition(20), it is clear that coproduct inclusions in $\mathcal{G}V$ are mono. Thus it suffices to show that f factors through a unique summand. Since X is non-empty it suffices to show that f sends any pair (a, b) of elements of X_0 to the same summand. Since $\pi_0(X) = 1$ there is a sequence

$$(x_j : 0 \leq j \leq n)$$

of elements of X_0 , such that $x_0 = a$, $x_n = b$ and for all $1 \leq j \leq n$, the set

$$S_{i,j} := V(X(x_{j-1}, x_j), \emptyset) \times V(X(x_j, x_{j-1}), \emptyset)$$

is empty. But if fx_{j-1} and fx_j are in different components of the coproduct, then both $Y(fx_{j-1}, fx_j)$ and $Y(fx_j, fx_{j-1})$ would be $\emptyset$, and so the hom maps $(f_{x_{j-1}, x_j}, f_{x_j, x_{j-1}})$ would give an element of $S_{i,j}$. Thus all the elements $(x_0, \dots, x_n)$ are sent to the same component of the coproduct by f .

Conversely suppose that X is connected. In general the $c_{X,i}$ induce a map

$$k_X : \coprod_{i \in \pi_0(X)} X_i \rightarrow X$$

which is the identity on objects. For any pair (a, b) in different components of X one can choose a map $r_{X,a,b} : X(a, b) \rightarrow \emptyset$ (using the axiom of choice). Together with the identities in the case of (a, b) in the same component, these provide the hom maps for a retraction r_X to k_X . Since X is connected r_X factors through a unique summand X_i , and so X_i is the only component of X whence $\pi_0(X) = 1$. $\square$

Despite this result, the notion of connected component only behaves completely as expected when $\emptyset \in V$ is a strict initial object.

Proposition 23. *Suppose that the category V has an initial object $\emptyset$. Then the following are equivalent:*

- (1) $\emptyset$ is a strict initial object.
- (2) For all $X \in \mathcal{G}V$, the maps $c_{X,i}$ exhibit X as a coproduct of its connected components.
- (3) $\mathcal{G}V$ is extensive.

PROOF. (1) $\Rightarrow$ (2): By strict initiality the hom maps $r_{X,a,b} : X(a, b) \rightarrow \emptyset$ are isomorphisms, and so r is fully faithful. However it is also the identity on objects, and thus r_X and k_X are (mutually inverse) isomorphisms.

(2) $\Rightarrow$ (3): Coproduct inclusions in $\mathcal{G}V$ are fully faithful by construction, and so by proposition(20), corollary(19) and (2), $\mathcal{G}V$ satisfies the hypotheses of lemma(2), from which the result follows.

(3) $\Rightarrow$ (1): Given a map $f : X \rightarrow \emptyset$ in V , one obtains an identity on objects map $(f) : (X) \rightarrow (\emptyset)$ in $\mathcal{G}V$. But $(\emptyset) = 0 + 0$, and so pulling back f along the coproduct inclusions one obtains, by the extensivity of $\mathcal{G}V$ a coproduct diagram

$$A \xrightarrow{c_A} (X) \xleftarrow{c_B} B$$

with A and B each having one object. Since $(-)_0$ preserves this coproduct by proposition(20), c_A and c_B pick out different objects, and so by the construction of coproducts in $\mathcal{G}V$, one has $X = (X)(0, 1) \cong \emptyset$. $\square$

Now we describe how the 2-functor $\mathcal{G}$ preserves locally (c)-presentable categories and Grothendieck toposes. First we require a general lemma which produces a strongly generating or dense subcategory of $\mathcal{G}V$ from one in V in a canonical way. Recall that an object D of a category V is said to be *small projective* when $V(D, -)$ preserves all small colimits.

Lemma 24. *Let $\mathcal{D}$ be a full subcategory of V and suppose that V has an initial object. Define an associated full subcategory $\mathcal{D}_+$ of $\mathcal{G}V$ as follows:*

- $0 \in \mathcal{D}_+$.
- $D \in \mathcal{D} \Rightarrow (D) \in \mathcal{D}_+$.

Let λ be a regular cardinal.

- (1) *If $\mathcal{D}$ is a strong generator then so is $\mathcal{D}_+$. [18]*
- (2) *If $\mathcal{D}$ is dense then so is $\mathcal{D}_+$.*
- (3) *If the objects of $\mathcal{D}$ are connected then so are those of $\mathcal{D}_+$.*
- (4) *If V has λ -filtered colimits and the objects of $\mathcal{D}$ are λ -presentable, then so are those of $\mathcal{D}_+$. [18]*
- (5) *If V has small colimits and the objects of $\mathcal{D}$ are small projective, then so are those of $\mathcal{D}_+$.*

PROOF. Let $f : X \rightarrow Y$ be in $\mathcal{G}V$. Suppose that $\mathcal{G}V(0, f)$ is a bijection, and that for all $D \in \mathcal{D}$, $\mathcal{G}V((D), f)$ are bijections. For (1) we must show that f is an isomorphism. To say that $\mathcal{G}V(0, f)$ is a bijection is to say that f is bijective on objects, and so it suffices to show that f is fully faithful. Let $a, b \in X_0$ and $D \in \mathcal{D}$. Note that the hom set $V(D, X(a, b))$ may be recovered as the pullback of the cospan

$$1 \xrightarrow{(a,b)} \mathcal{G}V(0, X) \times \mathcal{G}V(0, X) \xleftarrow{(\mathcal{G}V(i_0, X), \mathcal{G}V(i_1, X))} \mathcal{G}V((D), X)$$

where i_0 and i_1 pick out the objects 0 and 1 of (D) respectively. Moreover the function $V(D, f_{a,b})$ is induced by the isomorphism of cospans

$$\begin{array}{ccc} 1 & \xrightarrow{(a,b)} & \mathcal{G}V(0, X) \times \mathcal{G}V(0, X) \xleftarrow{(\mathcal{G}V(i_0, X), \mathcal{G}V(i_1, X))} \mathcal{G}V((D), X) \\ \downarrow & & \downarrow \mathcal{G}V(0, f) \times \mathcal{G}V(0, f) \qquad \qquad \downarrow \mathcal{G}V((D), f) \\ 1 & \xrightarrow{(fa,fb)} & \mathcal{G}V(0, Y) \times \mathcal{G}V(0, Y) \xleftarrow{(\mathcal{G}V(i_0, Y), \mathcal{G}V(i_1, Y))} \mathcal{G}V((D), Y) \end{array}$$

and so $V(D, f_{a,b})$ is also bijective. Since this is true for all $D \in \mathcal{D}$ and $\mathcal{D}$ is a strong generator, it follows that $f_{a,b}$ is an isomorphism. Thus f is fully faithful and so (1) follows.

Given functions

$$f_E : \mathcal{G}V(E, X) \rightarrow \mathcal{G}V(E, Y)$$

natural in $E \in \mathcal{D}_+$, we must show for (2) that there is a unique $f : X \rightarrow Y$ such that $f_E = \mathcal{G}V(E, f)$. The object map of f is forced to be f_0 , and naturality with respect to the maps i_0 and $i_1 : 0 \rightarrow (D)$ ensures that the functions f_E amount to f_0 together with functions

$$f_{D,a,b} : \mathcal{G}(t_V)^{\times 2}_{\bullet}((0, (D), 1), (a, X, b)) \rightarrow \mathcal{G}(t_V)^{\times 2}_{\bullet}((0, (D), 1), (f_0 a, Y, f_0 b))$$

natural in $D \in \mathcal{D}$ for all $a, b \in X_0$. Recall from section(3.1) that $\varepsilon_V : \mathcal{G}(t_V)^{\times 2}_{\bullet} \rightarrow V$ has a left adjoint given on objects by $Z \mapsto (0, (Z), 1)$. By this adjointness the above

maps are in turn in bijection with maps

$$f'_{D,a,b} : V(D, X(a, b)) \rightarrow V(D, Y(f_0 a, f_0 b))$$

natural in $D \in \mathcal{D}$ for all $a, b \in X_0$, and so by the density of $\mathcal{D}$ one has unique $f_{a,b}$ in V such that $f'_{D,a,b} = V(D, f_{a,b})$. Thus f_0 and the $f_{a,b}$ together form the object and hom maps of the unique desired map f , and so (2) follows.

By proposition(20) $(-)_0$ preserves all the necessary colimits, so that in the case of (3) 0 is connected, in the case of (4) it is λ -presentable and in the case of (5) it is small projective.

Recall the uniform construction of a colimit of $F : I \rightarrow \mathcal{G}V$ described for proposition(20), and write $\kappa_i : Fi \rightarrow K$ for the universal cocone. Then for $D \in V$ the cocone $\mathcal{G}V((D), \kappa_i)$ induces an obstruction map

$$\gamma_{(D), \kappa_i} : \operatorname{colim}_{i \in I} \mathcal{G}V((D), Fi) \rightarrow \mathcal{G}V((D), K)$$

which measures the extent to which $\mathcal{G}V((D), -)$ preserves the colimit of F . We shall give an alternative description of this map in terms of the homs of V . First observe that any map $f : (D) \rightarrow X$ amounts to an ordered pair (a, b) of objects of X picked out by f_0 , and the hom map $f_{0,1} : D \rightarrow X(a, b)$, and so one has a bijection

$$\mathcal{G}V((D), X) \cong \coprod_{a,b \in X_0} V(D, X(a, b)).$$

Second for $a, b \in K_0$ recall from the construction of the colimit K that one has a colimit cocone $\kappa_{i,\alpha,\beta} : Fi(\alpha, \beta) \rightarrow K(a, b)$ in V where (α, i, β) ranges over the fibre of $\kappa_{0\bullet}^{\times 2} : F_{0\bullet}^{\times 2} \rightarrow K_0 \times K_0$ over (a, b) . Thus one has an obstruction map

$$\gamma_{D, \kappa_{i,\alpha,\beta}} : \operatorname{colim}_{(i,\alpha,\beta) \in (\kappa_{0\bullet}^{\times 2})^{-1}(a,b)} V(D, Fi(\alpha, \beta)) \rightarrow V(D, K(a, b))$$

measuring the extent to which $V(D, -)$ preserves the defining colimit of $K(a, b)$. The above isomorphisms exhibit $\gamma_{(D), \kappa_i}$ as isomorphic in $\mathbf{Set}^{\rightarrow}$ to the function

$$\coprod_{a,b} \gamma_{D, \kappa_{i,a,b}} : \coprod_{a,b} \operatorname{colim}_{(\alpha,i,\beta)} V(D, Fi(\alpha, \beta)) \rightarrow \coprod_{a,b} V(D, K(a, b)).$$

Let D be small projective and I small. Then the colimit in the definition of $\gamma_{D, \kappa_{i,\alpha,\beta}}$ is preserved since D is small projective. Thus the functions $\gamma_{D, \kappa_{i,\alpha,\beta}}$, and thus $\gamma_{(D), \kappa_i}$ are bijective, whence (D) is also small projective, and so (5) follows. Similar arguments prove (4) and (3). In the case of (3) when D is connected and I discrete, $(\kappa_{0\bullet}^{\times 2})^{-1}(a, b)$ is either the empty or the terminal category. In the former case the colimit in the definition of $\gamma_{D, \kappa_{i,\alpha,\beta}}$ is preserved since D is connected, and in the latter case this is so since the colimit in question is absolute. As for (4) where D is now λ -presentable and I is λ -filtered, the result follows because as we saw in the proof of proposition(20), the categories $(\kappa_{0\bullet}^{\times 2})^{-1}(a, b)$ are also λ -filtered. $\square$

Theorem 25. *Let λ be a regular cardinal.*

- (1) *If V is locally λ -presentable then so is $\mathcal{G}V$. [18]*
- (2) *If V is locally λ -presentable and has a strict initial object, then $\mathcal{G}V$ is locally λ -c-presentable.*
- (3) *If V is locally λ -presentable then $\mathcal{G}^2 V$ is locally λ -c-presentable.*
- (4) *If V is locally λ -c-presentable then so is $\mathcal{G}V$.*
- (5) *If V is a presheaf topos then so is $\mathcal{G}V$.*

(6) *If V is a Grothendieck topos then $\mathcal{G}V$ is a locally connected Grothendieck topos.*

PROOF. If V is locally λ -presentable then $\mathcal{G}V$ is cocomplete by proposition(20), and one can build a strong generator for $\mathcal{G}V$ consisting of λ -presentable objects from one in V using lemma(24) to exhibit $\mathcal{G}V$ as locally λ -presentable, giving us (1). If in addition V has a strict initial object, then in $\mathcal{G}V$ every object decomposes as a sum connected objects by proposition(23), and so (2) follows by theorem(8)(8). (3) now follows since $\mathcal{G}V$ has a strict initial object by proposition(20). (4) is immediate from (2) and theorem(8)(9).

Recall that a category V is a presheaf topos iff it has a small dense full subcategory $i : \mathcal{D} \hookrightarrow V$ consisting of small projective objects. Clearly the representables in a presheaf topos provide such a dense subcategory. Conversely $V(i, 1) : V \rightarrow \widehat{\mathcal{D}}$ is fully faithful by density. Since the objects of $\mathcal{D}$ are small projective, $V(i, 1)$ is cocontinuous, and since every presheaf is a colimit of representables, it then follows that $V(i, 1)$ is essentially surjective on objects, giving the desired equivalence $V \simeq \widehat{\mathcal{D}}$. In this situation $\mathcal{D}_+$ provides a small dense subcategory of $\mathcal{G}V$ consisting of small projectives by lemma(24), whence $\mathcal{G}V \simeq \widehat{\mathcal{D}_+}$, and so (5) follows.

Since a Grothendieck topos is a left exact localisation of a presheaf category, the 2-functoriality of $\mathcal{G}$ together with (5), (2) and example(9) implies that to establish (6), it suffices to show that $\mathcal{G}$ preserves left exact functors between categories with finite limits. This follows immediately from the explicit description of limits in $\mathcal{G}V$ given in the proof of proposition(18). $\square$

For a small category $\mathbb{C}$, lemma(24) and theorem(25)(5) gives us a category $\mathbb{C}_+$ such that $\mathcal{G}\widehat{\mathbb{C}} \simeq \widehat{\mathbb{C}_+}$. A syntactic description $\mathbb{C}_+$ is as follows. There is an injective on objects fully faithful functor

$$i_{\mathbb{C}} : \mathbb{C} \rightarrow \mathbb{C}_+ \quad C \mapsto C_+$$

and $\mathbb{C}_+$ has an additional object 0 not in the image of $i_{\mathbb{C}}$. Moreover for each $C \in \mathbb{C}$ one has maps

$$\sigma_C : 0 \rightarrow C_+ \quad \tau_C : 0 \rightarrow C_+$$

and for all $f : C \rightarrow D$ one has the equations $f_+ \sigma_C = \sigma_D$ and $f_+ \tau_C = \tau_D$. Starting from the terminal category and iterating $(-)_+$ n times gives the usual site

$$0 \begin{array}{c} \xrightarrow{\sigma} \\ \xrightarrow{\tau} \end{array} \dots \begin{array}{c} \xrightarrow{\sigma} \\ \xrightarrow{\tau} \end{array} n$$

for n -globular sets.

4. The multitensor to monad construction

4.1. Recalling multitensors. We begin by recalling some definitions and notation from [6]. For a category V , the free strict monoidal category MV on V is described as follows. An object of MV is a finite sequence $(Z_1, \dots, Z_n)$ of objects of V . A map is a sequence of maps of V – there are no maps between sequences of objects of different lengths. The unit $\eta_V : V \rightarrow MV$ of the 2-monad M is the inclusion of sequences of length 1. The multiplication $\mu_V : M^2V \rightarrow MV$ is given by concatenation. A *lax monoidal category* is a lax algebra for the 2-monad M .

Explicitly a multitensor on a category V consists of a functor $E : MV \rightarrow V$, and maps

$$u_Z : Z \rightarrow E(Z) \quad \sigma_{Z_{ij}} : E_i E_j Z_{ij} \rightarrow E_{ij} Z_{ij}$$

for all Z, Z_{ij} from V which are natural in their arguments, and such that

$$\begin{array}{ccccc} E_i Z_i \xrightarrow{u_i} E_1 E_i Z_i & E_i E_j E_k Z_{ijk} \xrightarrow{\sigma_k} E_{ij} E_k Z_{ijk} & E_i E_1 Z_i \xleftarrow{E_i u} E_i Z_i \\ \downarrow 1 & \downarrow E_i \sigma & \searrow \sigma & \downarrow 1 \\ E_i Z_i & E_i E_{jk} Z_{ijk} \xrightarrow{\sigma} E_{ijk} Z_{ijk} & E_i Z_i \end{array}$$

in V . As in [6] the expressions

$$E(X_1, \dots, X_n) \quad \bigcup_{1 \leq i \leq n} E_i X_i \quad E_i X_i$$

are alternative notation for the n -ary tensor product of the objects $X_1, \dots, X_n$, and we refer to the endofunctor of V obtained by observing the effect of E on singleton sequences as E_1 . The data (E, u, σ) is called a *multitensor* on V , and u and σ are referred to as the *unit* and *substitution* of the multitensor respectively.

Given a multitensor (E, u, σ) on V , a *category enriched in E* consists of $X \in \mathcal{GV}$ together with maps

$$\kappa_{x_i} : E_i X(x_{i-1}, x_i) \rightarrow X(x_0, x_n)$$

for all $n \in \mathbb{N}$ and sequences $(x_0, \dots, x_n)$ of objects of X , such that

$$\begin{array}{ccc} X(x_0, x_1) \xrightarrow{u} E_1 X(x_0, x_1) & E_i E_j X(x_{(ij)-1}, x_{ij}) \xrightarrow{\sigma} E_{ij} X(x_{(ij)-1}, x_{ij}) \\ \searrow \text{id} & \downarrow E_i \kappa & \downarrow \kappa \\ & X(x_0, x_1) & E_i X(x_{(i1)-1}, x_{in_i}) \xrightarrow{\kappa} X(x_0, x_{mn_m}) \end{array}$$

commute, where $1 \leq i \leq m$, $1 \leq j \leq n_i$ and $x_{(11)-1} = x_0$. Since a choice of i and j references an element of the ordinal $n_\bullet$, the predecessor $(ij)-1$ of the pair (ij) is well-defined when i and j are not both 1. With the obvious notion of E -functor (see [6]), one has a category $E\text{-Cat}$ of E -categories and E -functors together with a forgetful functor

$$U^E : E\text{-Cat} \rightarrow \mathcal{GV}.$$

A multitensor (E, u, σ) is *distributive* when for all n the functor

$$V^n \rightarrow V \quad (X_1, \dots, X_n) \mapsto E(X_1, \dots, X_n)$$

preserves coproducts in each variable.

Multitensors generalise non-symmetric operads. For given a non-symmetric operad

$$(A_n : n \in \mathbb{N}) \quad u : I \rightarrow A_1 \quad \sigma : A_k \otimes A_{n_1} \otimes \dots \otimes A_{n_k} \rightarrow A_n.$$

in a braided monoidal category V , one can define a multitensor E on V via the formula

$$\bigcup_{1 \leq i \leq n} E_i X_i = A_n \otimes X_1 \otimes \dots \otimes X_n$$

as observed in [6] example(2.6), and when the tensor product for V is distributive, so is E . A category enriched in E with one object is precisely an A -algebra in the usual sense. In the case where V is **Set** with tensor product given by cartesian product, this construction is part of an equivalence between the evident category of distributive multitensors on **Set** and that of non-symmetric operads in **Set**. This equivalence is easily established using the fact that every set is a coproduct of singletons.

4.2. Monads from multitensors. Let (E, u, σ) be a distributive multitensor on a category V with coproducts. Then we define a monad ΓE on the category $\mathcal{GV}$ of graphs enriched in V as follows. We ask that the monad ΓE actually live over **Set**, that is to say, in the 2-category **CAT/Set**. Thus for $X \in \mathcal{GV}$, $\Gamma E(X)$ has the same objects as X . The homs of $\Gamma E(X)$ are defined by the equation

$$(2) \quad \Gamma E(X)(a, b) = \coprod_{a=x_0, \dots, x_n=b} \coprod_i E X(x_{i-1}, x_i)$$

for all $a, b \in X_0$. The above coproduct is taken over all finite sequences of objects of X starting at a and finishing at b . Let us write $k_{E, X, (x_i)_i}$ for a given coproduct inclusion for the above sum. Since the monad we are defining is over **Set**, the object maps of the components of the unit η and multiplication μ are identities, and so it suffices to define their hom maps. For the unit these are the composites

$$X(a, b) \xrightarrow{u_{X(a, b)}} E_1 X(a, b) \xrightarrow{k_{E, X, a, b}} \Gamma E(X)(a, b).$$

In order to define the multiplication, observe that the composites

$$\coprod_i \coprod_j E X(x_{ij-1}, x_{ij}) \xrightarrow{\coprod_i k} \coprod_i E \Gamma E(X)(x_{i-1}, x_i) \xrightarrow{k} (\Gamma E)^2(X)(a, b)$$

ranging of all doubly-nested sequences $(x_{ij})_{ij}$ of objects of X starting from a and finishing at b , exhibit the hom $(\Gamma E)^2(X)(a, b)$ as a coproduct, since E preserves coproducts in each variable. Let us write $k_{E, X, (x_{ij})_{ij}}^{(2)}$ for such a coproduct inclusion. We can now define the hom map of the components of the multiplication μ as unique such that

$$\begin{array}{ccc} \coprod_i \coprod_j E X(x_{ij-1}, x_{ij}) & \xrightarrow{\sigma} & \coprod_{ij} E X(x_{ij-1}, x_{ij}) \\ k^{(2)} \downarrow & & \downarrow k \\ (\Gamma E)^2(X)(a, b) & \xrightarrow{\mu_{X, a, b}} & \Gamma E(X)(a, b) \end{array}$$

commutes for all doubly-nested sequences $(x_{ij})_{ij}$ starting at a and finishing at b .

Proposition 26. *Let V be a category with coproducts and (E, u, σ) be a distributive multitensor on V . Then $(\Gamma E, \eta, \mu)$ as defined above is a monad on $\mathcal{GV}$ and one has an isomorphism $E\text{-Cat} \cong (\mathcal{GV})^{\Gamma E}$ commuting with the forgetful functors into $\mathcal{GV}$.*

PROOF. Since $(\Gamma E, \eta, \mu)$ are defined over **Set** it suffices to check the monad axioms on the homs. For the unit laws we must verify the commutativity of

$$\begin{array}{ccc} \Gamma E(X)(a, b) & \xrightarrow{\eta_{\Gamma E}} & (\Gamma E)^2(X)(a, b) \\ & \searrow 1 & \downarrow \mu \\ & & \Gamma E(X)(a, b) \end{array} \quad \begin{array}{ccc} (\Gamma E)^2(X)(a, b) & \xleftarrow{\Gamma E(\eta)} & \Gamma E(X)(a, b) \\ & \swarrow \mu & \nwarrow 1 \\ & & \Gamma E(X)(a, b) \end{array}$$

and precomposing each of these by each of the injections $k_{E,X,(x_i)_i}$ gives the unit laws for the multitensor E . Given a triply-nested sequence of objects of X starting at a and finishing at b , let us denote by $k_{E,X,(x_{ijk})_{ijk}}^{(3)}$ the composite

$$E_i E_j E_k X(x_{ijk-1}, x_{ijk}) \xrightarrow{E_i k^{(2)}} E_i(\Gamma E)^2(X)(x_{i-1}, x_i) \xrightarrow{k} (\Gamma E)^3(X)(a, b)$$

and note that since E is distributive, the family of maps so determined exhibits the hom $(\Gamma E)^3(X)(a, b)$ as a coproduct. The associative law on the homs then follows because precomposing the diagrams that express it with such coproduct injections gives back the associativity diagrams for the multitensor E . Thus $(\Gamma E, \eta, \mu)$ is indeed a monad on $\mathcal{GV}$.

For $X \in \mathcal{GV}$ a morphism $a : \Gamma E(X) \rightarrow X$ may be identified, by precomposing with the appropriate coproduct inclusions, with morphisms $E_i X(x_{i-1}, x_i) \rightarrow X(x_0, x_n)$ for all sequences $(x_0, \dots, x_n)$ of objects of X , and under this identification the unit and associative laws for a ΓE -algebra correspond exactly to those for an E -category. To say that $f : X \rightarrow Y$ in $\mathcal{GV}$ underlies a given morphism $(X, a) \rightarrow (Y, b)$ of ΓE -algebras is clearly equivalent to saying that f underlies an E -functor. Thus one has the required canonical isomorphism over $\mathcal{GV}$. $\square$

Example 27. In the case where V is **Set** and E is cartesian product, ΓE is the monad for categories on **Gph**. The summand of equation(2) corresponding to a given sequence $(x_0, \dots, x_n)$ is the set of paths in X of length n , starting at $a = x_0$ and finishing at $b = x_n$, which visits successively the intermediate vertices $(x_1, \dots, x_{n-1})$.

Remark 28. Given a monad T on $\mathcal{GV}$ over **Set**, and a set Z , one obtains by restriction a monad T_Z on the category $\mathcal{GV}_Z$ of V -graphs with fixed object set Z . Let us write Γ^{old} for the functor labelled as Γ in [6]. Then for a given distributive multitensor E , our present Γ and Γ^{old} are related by the formula

$$\Gamma^{\text{old}}(E) = \Gamma(E)_1$$

where the 1 on the right hand side of this equation indicates a singleton. In other words in this paper we are describing the “many-objects version” of the theory presented in [6] section(4).

4.3. Properties of ΓE . For a functor

$$F : A_1 \times \dots \times A_n \rightarrow B$$

of n variables, the preservation by F of a given connected limit or colimit implies that this limit or colimit is preserved in each variable separately. To see this one considers diagrams which are constant in all but the variable of interest, and use the fact that the limit/colimit of a connected diagram constant at an object X , is X , as witnessed by a universal cone/cocone all of whose components are 1_X .

However the converse of this is false in general. For instance to say that F preserves pullbacks is to say that it does so in each variable, and moreover, that all

squares of the form

(3)

$$\begin{array}{ccc} (a_1, \dots, a_{i-1}, a_i, \dots, a_j, a_{j+1}, \dots, a_n) & \xrightarrow{(1, \dots, 1, f, 1, \dots, 1)} & (a_1, \dots, a_{i-1}, b_i, \dots, a_j, a_{j+1}, \dots, a_n) \\ \downarrow (1, \dots, 1, g, 1, \dots, 1) & & \downarrow (1, \dots, 1, g, 1, \dots, 1) \\ (a_1, \dots, a_{i-1}, a_i, \dots, b_j, a_{j+1}, \dots, a_n) & \xrightarrow{(1, \dots, 1, f, 1, \dots, 1)} & (a_1, \dots, a_{i-1}, b_i, \dots, b_j, a_{j+1}, \dots, a_n) \end{array}$$

are sent to pullbacks in B , where $1 \leq i < j \leq n$, $f : a_i \rightarrow b_i$ and $g : a_j \rightarrow b_j$. That this extra condition follows from F preserving all pullbacks follows since these squares are obviously pullbacks in $A_1 \times \dots \times A_n$. Conversely note that any general map $(f_1, \dots, f_n) : (a_1, \dots, a_n) \rightarrow (b_1, \dots, b_n)$ in $A_1 \times \dots \times A_n$ can be factored in the following manner

$$(a_1, \dots, a_n) \xrightarrow{(f_1, 1, \dots, 1)} (b_1, a_2, \dots, a_n) \xrightarrow{(1, f_2, \dots, 1)} \dots \xrightarrow{(1, \dots, 1, f_n)} (b_1, \dots, b_n)$$

and doing so to each of the maps in a general pullback in $A_1 \times \dots \times A_n$, produces an $n \times n$ lattice diagram in which each inner square is either of the form (3), or a pullback in a single variable.

An important case where such distinctions can be ignored is with λ -filtered colimits for some regular cardinal λ . For suppose that F preserves λ -filtered colimits in each variable. By [2] corollary(1.7) it suffices to show that F preserves colimits of chains of length λ . Given such a chain

$$X : \lambda \rightarrow A_1 \times \dots \times A_n \quad i \mapsto (X_{i1}, \dots, X_{in})$$

with object map denoted on the right, one obtains the functor

$$X' : \lambda^n \rightarrow A_1 \times \dots \times A_n \quad (i_1, \dots, i_n) \mapsto (X_{i_1 1}, \dots, X_{i_n n})$$

which one may readily verify has the same colimit as X . But the colimit of X' may be taken one variable at a time and so

$$\text{colim}(X) \cong \text{colim}_{i_1} \text{colim}_{i_2} \dots \text{colim}_{i_n} (X_{i_1 1}, \dots, X_{i_n n})$$

from which it follows that F preserves $\text{colim}(X)$. We say that a multitensor (E, u, σ) is λ -*accessible* when the functor $E : MV \rightarrow V$ preserves λ -filtered colimits, which is clearly equivalent to the condition that each of the associated n -ary functors $E_n : V^n \rightarrow V$ does so, which as we have seen, is equivalent to the condition that each of the E_n 's preserve λ -filtered colimits in each variable.

Cartesian monads play a fundamental role in higher category theory [21]. Recall that a monad (T, η, μ) on a category V with pullbacks is said to be *cartesian* when T preserves pullbacks, and η and μ are cartesian transformations (meaning that their naturality squares are pullbacks). Similarly one has the notion of a cartesian multitensor, with a multitensor (E, u, σ) on a category V with pullbacks being *cartesian* when E preserves pullbacks, and u and σ are cartesian transformations.

Recall that a functor $F : V \rightarrow W$ is a *local right adjoint* (l.r.a) when for all $X \in V$ the induced functor

$$F_X : V/X \rightarrow V/FX$$

between slice categories is a right adjoint. When V has a terminal object 1, it suffices for l.r.a-ness that F_1 be a right adjoint. Recall moreover that l.r.a functors preserve all connected limits, and thus in particular pullbacks. A monad (T, η, μ)

on a category V is l.r.a (as a monad) when T is l.r.a as a functor and η and μ are cartesian. Thus this is a slightly stronger condition on a monad than being cartesian. Local right adjoint monads, especially defined on presheaf categories, are fundamental to higher category theory. Indeed a deeper understanding of such monads is the key to understanding the relationship between the operadic and homotopical approaches to the subject [26]. Similarly one has the notion of an l.r.a multitensor, with a multitensor (E, u, σ) on a category V being l.r.a when the functor $E : MV \rightarrow V$ is l.r.a, and u and σ are cartesian transformations.

For a functor $F : \coprod_{i \in I} V_i \rightarrow W$ to be l.r.a is equivalent to each of the induced $F_i : V_i \rightarrow W$ being l.r.a, because for $X \in V_i$, $F_X = (F_i)_X$. Thus the condition that $E : MV \rightarrow V$ be l.r.a is equivalent to the condition that each of the E_n 's is l.r.a. The condition that a functor $F : V_1 \times \dots \times V_n \rightarrow W$ to be l.r.a is equivalent to the condition that it be l.r.a in each variable, and moreover that it send the basic pullbacks (3) in $V_1 \times \dots \times V_n$ to pullbacks in W . For suppose F is l.r.a. Then since it preserves all pullbacks it preserves those of the form (3). Moreover for $1 \leq i \leq n$ the functor

$$F(X_1, \dots, X_{i-1}, -, X_{i+1}, \dots, X_n)_{X_i} : V_i/X_i \rightarrow W/F(X_1, \dots, X_n)$$

can be written as the composite

$$V_i/X_i \longrightarrow V_1/X_1 \times \dots \times X_n/X_n \xrightarrow{F_{X_1, \dots, X_n}} W/F(X_1, \dots, X_n)$$

in which the first functor has object map $f \mapsto F(1_{X_1}, \dots, 1_{X_{i-1}}, f, 1_{X_{i+1}}, \dots, 1_{X_n})$. Since for all i both these functors are clearly right adjoints, F is l.r.a in each variable. Conversely, supposing F to be l.r.a in each variable and preserving the pullbacks (3), F 's effect on the slice over $(X_1, \dots, X_n)$ is isomorphic to the composite

$$\prod_i V/X_i \xrightarrow{\prod_i F(X_1, \dots, X_{i-1}, -, X_{i+1}, \dots, X_n)_{X_i}} (W/F(X_1, \dots, X_n))^n \xrightarrow{\times} W/F(X_1, \dots, X_n)$$

and both these functors are clearly right adjoints. Thus the condition that $E : MV \rightarrow V$ be l.r.a is equivalent to the condition that each E_n is l.r.a in each variable and preserve the pullbacks of the form (3).

The following result expresses how the assignment $E \mapsto \Gamma E$ is compatible with the various categorical properties we have been discussing.

Theorem 29. *Let V be a category with coproducts and (E, u, σ) be a distributive multitensor on V , and let $(\Gamma E, \eta, \mu)$ be the corresponding monad on $\mathcal{G}V$. Let λ be a regular cardinal.*

- (1) ΓE preserves coproducts.
- (2) Suppose V has λ -filtered colimits. Then E is λ -accessible iff ΓE is.
- (3) Suppose V has pullbacks and every object of V is a coproduct of connected objects. Then (E, u, σ) is a cartesian multitensor iff $(\Gamma E, \eta, \mu)$ is a cartesian monad.
- (4) Suppose V is locally λ -c-presentable. Then (E, u, σ) is an l.r.a multitensor iff $(\Gamma E, \eta, \mu)$ is an l.r.a monad.

The proof of this result will occupy the remainder of section(4).

4.4. Coproducts and filtered colimits. In lemma(30) below we formulate the preservation by ΓE of a given colimit in terms of the underlying multitensor E . We require some further notation. For a functor $f : J \rightarrow \mathbf{Set}$ and $n \in \mathbb{N}$ we denote

by $f^{\times n} : J \rightarrow \mathbf{Set}$ the functor with object map $j \mapsto f(j)^n$, and if $\kappa_j : fj \rightarrow K$ form a colimit cocone, then we denote by $\kappa_{\bullet}^{\times n} : f_{\bullet}^{\times n} \rightarrow K^n$ the evident induced functor. We have been using this notation already, for instance in proposition(20), in the case $n = 2$.

Lemma 30. *Let J be a small category, $F : J \rightarrow \mathcal{GV}$ and V has sufficient colimits so that the colimit K of F may be constructed as in the discussion preceeding proposition(20). Let $\kappa_{j,0} : F(j)_0 \rightarrow K_0$ be a colimit cocone in $\mathbf{Set}$ at the level of objects, and for $a, b \in K_0$ let*

$$\kappa_{j,\alpha,\beta} : F(j)(\alpha, \beta) \rightarrow K(a, b)$$

be the colimit cocone in V , where $(j, \alpha, \beta) \in (\kappa_{0\bullet}^{\times 2})^{-1}(a, b)$. If for all sequences $(x_0, \dots, x_n)$ of objects of K , the morphisms

$$\mathbb{E}_i \kappa_{j, \gamma_{i-1}, \gamma_i} : \mathbb{E}_i F(j)(\gamma_{i-1}, \gamma_i) \rightarrow \mathbb{E}_i K(x_{i-1}, x_i)$$

ranging over $(j, \gamma_0, \dots, \gamma_n) \in (\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n)$ form a colimit cocone in V , then ΓE preserves the colimit of F .

PROOF. The obstruction map k measuring whether ΓE preserves the colimit K is bijective on objects since ΓE is over $\mathbf{Set}$. By definition of ΓE and the construction of colimits in $\mathcal{GV}$ one has

$$\left(\text{colim}_{j \in J} \Gamma E(Fj) \right) (a, b) = \coprod_{a=x_0, \dots, x_n=b} \text{colim}_{j, \gamma_0, \dots, \gamma_n} \mathbb{E}_i F(j)(\gamma_{i-1}, \gamma_i)$$

where in the summand $(j, \gamma_0, \dots, \gamma_n) \in (\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n)$. Thus if the obstruction maps measuring whether the $\mathbb{E}_i \kappa_{j, \gamma_{i-1}, \gamma_i}$ are colimit cocones are invertible, then the hom maps of k are invertible, and so k is also fully faithful. $\square$

In order to understand how the preservation by E of λ -filtered colimits gives rise to the same property for ΓE , we require

Lemma 31. *Let J be a filtered category, $F : J \rightarrow \mathbf{Set}$ and $\kappa_j : F(j) \rightarrow K$ be a colimit cocone. Then for $n > 0$ and $1 \leq i \leq n$ the functor*

$$\text{pr}_i : (\kappa_{\bullet}^{\times n})^{-1}(x_0, \dots, x_n) \rightarrow (\kappa_{\bullet}^{\times 2})^{-1}(x_{i-1}, x_i) \quad (j, \gamma_0, \dots, \gamma_n) \mapsto (j, \gamma_{i-1}, \gamma_i)$$

is final.

PROOF. For a given $(j, \alpha, \beta) \in (\kappa_{\bullet}^{\times n})^{-1}(x_{i-1}, x_i)$ we must show that the comma category $(j, \alpha, \beta)/\text{pr}_i$ is connected. Explicitly the objects of this comma category consist of the data

$$f : j \rightarrow j' \quad (j', \gamma_0, \dots, \gamma_n)$$

where $\gamma_i \in Fj$, $F(f)(\alpha) = \gamma_{i-1}$ and $F(f)(\beta) = \gamma_i$. A morphism

$$(f, j', \gamma_0, \dots, \gamma_n) \rightarrow (f', j'', \gamma'_0, \dots, \gamma'_n)$$

is a map $g : j' \rightarrow j''$ in J such that $gf = f'$ and $F(g)(\gamma_k) = \gamma'_k$ for $1 \leq k \leq n$.

For $k \notin \{i-1, i\}$ one can find (j_k, γ_k) where $j_k \in J$, $\gamma_k \in F(j_k)$ and $\kappa_{j_k}(\gamma_k) = x_k$ since the cocone κ is jointly epic. By the filteredness of J one has maps $\delta : j \rightarrow j'$ and $\delta_k : j_k \rightarrow j'$, and thus $(\delta, \varepsilon_0, \dots, \varepsilon_n)$ with $\varepsilon_{i-1} = F(\delta)(\alpha)$, $\varepsilon_i = F(\delta)(\beta)$ and $\varepsilon_k = F(\delta)(\gamma_k)$ for $k \notin \{i-1, i\}$, exhibits $(j, \alpha, \beta)/\text{pr}_i$ as non-empty.

Note that if $y, z \in Fj$ satisfy $\kappa_j(x) = \kappa_j(y)$, then since K may be identified as the connected components of $F_\bullet$, there is an undirected path

$$(j, x) \rightarrow (j_1, z_1) \leftarrow \dots \rightarrow (j_n, z_n) \rightarrow (j, y)$$

in $F_\bullet$. Consider the underlying diagram in J with endpoints (ie the two instances of j) identified. Using the filteredness of J one has a cocone for this diagram, and we write j' for the vertex of this cocone. Thus we have $f : j \rightarrow j'$ such that $F(f)(y) = F(f)(z)$.

Now let $(f, j', \gamma_0, \dots, \gamma_n)$ and $(f', j'', \gamma'_0, \dots, \gamma'_n)$ be any two objects of $(j, \alpha, \beta)/\text{pr}_i$. First we use the filteredness of J to produce a commutative square

$$\begin{array}{ccc} j & \xrightarrow{f} & j' \\ f' \downarrow & & \downarrow h_1 \\ j'' & \xrightarrow{g_1} & v_1 \end{array}$$

whose diagonal we denote as d_1 . Note that by definition $F(h_1)(\gamma_{i-1}) = F(d_1)(\alpha) = F(g_1)(\gamma'_{i-1})$ and $F(h_1)(\gamma_i) = F(d_1)(\beta) = F(g_1)(\gamma'_i)$, but we have no reason to suppose that $F(h_1)(\gamma_k) = F(g_1)(\gamma'_k)$ for $k \notin \{i-1, i\}$. However $F(h_1)(\gamma_k)$ and $F(g_1)(\gamma_k)$ are by definition identified by κ_{v_1} . Choosing one value of k and using the observation of the previous paragraph, we can find $r_1 : v_1 \rightarrow v_2$ such that $F(h_2)(\gamma_k) = F(g_2)(\gamma'_k)$ where $h_2 = r_1 h_1$ and $g_2 = r_1 g_1$. Do the same successively for all other $k \notin \{i-1, i\}$, so that in the end one has $h : j' \rightarrow v$ and $g : j'' \rightarrow v$ such that $hf = gf'$ whose common value we denote as d , and $F(h)(\gamma_k) = F(g)(\gamma'_k)$ for all $1 \leq k \leq n$. Denote by $\psi_k \in F(v)$ for the common value of $F(h)(\gamma_k) = F(g)(\gamma'_k)$. Thus one has

$$(f, j', \gamma_0, \dots, \gamma_n) \xrightarrow{h} (d, v, \psi_1, \dots, \psi_k) \xleftarrow{g} (f', j'', \gamma'_0, \dots, \gamma'_n)$$

in $(j, \alpha, \beta)/\text{pr}_i$. Thus $(j, \alpha, \beta)/\text{pr}_i$ is indeed connected. $\square$

With these preliminary results in hand we can now proceed to

PROOF. (of theorem(29)(1) and (2))

(1): Let J be small and discrete and $F : J \rightarrow \mathcal{G}V$. In the situation of lemma(30) with a given sequence $(x_0, \dots, x_n)$ from K_0 , if that sequence contains elements from different $F(j)$'s then the category $(\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n)$ will be empty, but by distributivity in this case $\text{E}_i K(x_{i-1}, x_i)$ will also be initial. On the other hand when the x_i all come from the same $F(j)$, one has

$$(\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n) = \prod_{1 \leq i \leq n} \kappa_{j0}^{-1}(x_i)$$

and then the universality of the cocone $\text{E}_i \kappa_{j, \gamma_{i-1}, \gamma_i}$ follows again from the distributivity of E .

(2): Suppose E is λ -accessible. Let J be λ -filtered, $F : J \rightarrow \mathcal{G}V$ and $(x_0, \dots, x_n)$ be a sequence from K_0 , where as in lemma(30), K is the colimit of F . Then one has a functor

$$(\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n) \rightarrow V^n \quad (j, \gamma_0, \dots, \gamma_n) \mapsto (F(j)(\gamma_{i-1}, \gamma_i))_{1 \leq i \leq n}$$

and we claim that

$$(\kappa_{j, \gamma_{i-1}, \gamma_i} : F(j)(\gamma_{i-1}, \gamma_i) \rightarrow K(x_{i-1}, x_i) \quad : \quad 1 \leq i \leq n)$$

is a colimit cocone in V^n for this functor. In the i -th variable $\kappa_{j,\gamma_{i-1},\gamma_i}$ is a cocone for the composite functor

$$(\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n) \xrightarrow{\text{pr}_i} (\kappa_{0\bullet}^{\times 2})^{-1}(x_{i-1}, x_i) \longrightarrow V$$

in which the second leg has colimit cocone given by the components $\kappa_{j,\gamma_{i-1},\gamma_i}$. Since pr_i is final by lemma(31), the cocone $(\kappa_{j,\gamma_{i-1},\gamma_i} : 1 \leq i \leq n)$ is indeed universal as claimed.

Now the category $(F_{0\bullet}^{\times n})$ comes with a discrete opfibration into J , and so its connected components are λ -filtered. But since λ -filtered colimits commute with finite products in **Set**, these connected components are exactly the fibres of $(\kappa_{0\bullet}^{\times n})$, and so for each sequence $(x_0, \dots, x_n)$, $(\kappa_{0\bullet}^{\times n})^{-1}(x_0, \dots, x_n)$ is λ -filtered. Thus by lemma(30) ΓE is λ -accessible.

Conversely suppose that ΓE is λ -accessible. For $F : J \rightarrow V$ with J where is λ -filtered, with colimit cocone $\kappa_j : Fj \rightarrow K$ we must show that the induced cocone

$$(4) \quad E(X_1, \dots, X_{i-1}, Fj, X_{i+1}, \dots, X_n) \rightarrow E(X_1, \dots, X_{i-1}, K, X_{i+1}, \dots, X_n)$$

is universal, for all $N \in \mathbb{N}$, $1 \leq i \leq n$ and $X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n \in V$. By remark(21) the cocone

$$(5) \quad (X_1, \dots, X_{i-1}, Fj, X_{i+1}, \dots, X_n) \rightarrow (X_1, \dots, X_{i-1}, K, X_{i+1}, \dots, X_n)$$

in $\mathcal{G}V$ is universal, and moreover that for any sequence $(Y_1, \dots, Y_n)$ of objects of V and $1 \leq a, b \leq n$ one has

$$\begin{aligned} \Gamma E(Y_1, \dots, Y_n)(0, n) &= \coprod_{0=x_0, \dots, x_n=n} E_i((Y_1, \dots, Y_n)(x_{i-1}, x_i)) \\ &\cong E_i Y_i \end{aligned}$$

by the distributivity of E . Thus applying ΓE to the cocone (5) and looking at the hom between 0 and n gives the cocone (4), and so by remark(21), the result follows. $\square$

4.5. Cartesianness of ΓE . Let V be a category with coproducts and pullbacks, in which every object is a coproduct of connected objects, and suppose that (E, u, σ) is a cartesian multitensor. We will now show that $(\Gamma E, \eta, \mu)$ is a cartesian monad. Note that by lemma(2) such a V is in fact extensive.

Lemma 32. *Let V be a category with coproducts and pullbacks in which every object is a coproduct of connected objects. Suppose that we are given square*

$$\begin{array}{ccc} P & \xrightarrow{q} & B \\ p \downarrow & & \downarrow g \\ A & \xrightarrow{f} & C \end{array}$$

in V which admits a description as on the left in

$$\begin{array}{ccc} \coprod_{(i,j) \in L} P_{ij} & \xrightarrow{(q_{ij})_{ij}} & \coprod_{j \in J} B_j \\ (p_{ij})_{ij} \downarrow & & \downarrow (g_j)_j \\ \coprod_{i \in I} A_i & \xrightarrow{(f_i)_i} & \coprod_{k \in K} C_k \end{array} \quad \begin{array}{ccc} L & \xrightarrow{\nu} & J \\ \pi \downarrow & pb & \downarrow \gamma \\ I & \xrightarrow{\phi} & K \end{array} \quad \begin{array}{ccc} P_{(i,j)} & \xrightarrow{q_{ij}} & B_j \\ p_{ij} \downarrow & & \downarrow g_j \\ A_i & \xrightarrow{f_i} & C_{\phi i = \gamma j} \end{array}$$

in which the indexing sets of the coproduct decompositions fit into a pullback square as shown in the middle, with elements of L represented explicitly as pairs (i, j) such that $\phi(i) = \gamma(j)$. Suppose moreover that for all such (i, j) the squares as indicated on the right in the previous display are pullbacks. Then it follows that the original square is itself a pullback.

PROOF. To see this is a pullback it suffices just for connected X , $h : X \rightarrow A$ and $k : X \rightarrow B$ with $fh = gk$, that there is a unique filler $d : X \rightarrow P$ such that $pd = h$ and $qd = k$, since every object of V is a coproduct of connected ones. But then using the connectedness one can factor h and k through unique summands say $i \in I$ and $j \in J$ related by $\phi(i) = \gamma(j)$, and so use the defining pullback of P_{ij} to induce the desired unique d . $\square$

One application of lemma(32) is the componentwise construction of pullbacks in such a V . For given a cospan

$$A \xrightarrow{f} C \xleftarrow{g} B$$

in V , one can compute its pullback one component at a time by decomposing A , B and C into coproducts of connected objects, then pulling back the indexing sets, then taking the pullbacks componentwise, and finally re-amalgamating (by taking coproducts). Note however that the summands P_{ij} of the pullback so obtained are not necessarily themselves connected. We are now ready to exhibit

PROOF. (of theorem(29)(3))

Let (E, u, σ) a cartesian multitensor on V a category with coproducts and pullbacks in which every object decomposes as sum of connected ones. Let $\mathbb{P}$ be the pullback

$$\begin{array}{ccc} P & \xrightarrow{q} & Y \\ p \downarrow & pb & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

in $\mathcal{G}V$ and denote by $d : P \rightarrow Z$ the diagonal. Then $\Gamma E(\mathbb{P})$ is certainly a pullback at the level of object sets, since ΓE is over **Set**. So it suffices, by the construction of pullbacks in $\mathcal{G}V$, to check that for each $w, w' \in P_0$ the corresponding hom square of $\Gamma E(\mathbb{P})$ is a pullback in V . This hom square is a square in V of the form

$$\begin{array}{ccc} \coprod_{w=w_0, \dots, w_n=w'} \mathbf{E}_i P(w_{i-1}, w_i) & \xrightarrow{\quad} & \coprod_{qw=y_0, \dots, y_n=qw'} \mathbf{E}_i Y(y_{i-1}, y_i) \\ \downarrow & & \downarrow \\ \coprod_{pw=x_0, \dots, x_n=pw'} \mathbf{E}_i X(x_{i-1}, x_i) & \xrightarrow{\quad} & \coprod_{dw=z_0, \dots, z_n=dw'} \mathbf{E}_i Z(z_{i-1}, z_i) \end{array}$$

and the induced square at the level of summand indexing sets is a pullback since $\mathbb{P}_0$ is a pullback in **Set**. For each sequence $(w_0, \dots, w_n)$ in P_0 from w to w' , the

corresponding component is

$$\begin{array}{ccc}
 E_i P(w_{i-1}, w_i) & \xrightarrow{E_i q_{w_{i-1}, w_i}} & E_i Y(qw_{i-1}, qw_i) \\
 E_i p_{w_{i-1}, w_i} \downarrow & & \downarrow E_i g_{qw_{i-1}, qw_i} \\
 E_i X(pw_{i-1}, pw_i) & \xrightarrow{E_i f_{pw_{i-1}, pw_i}} & E_i Z(dw_{i-1}, dw_i)
 \end{array}$$

which is a pullback since $\mathbb{P}$ is. Thus by lemma(32) $\Gamma E(\mathbb{P})$ is a pullback.

We must show that for $f : X \rightarrow Y$ in $\mathcal{GV}$ the corresponding naturality squares of η and μ are cartesian. Since they are over **Set** this is clearly so at the level of objects. The hom at (a, b) of the naturality of square of η has underlying square of summand indexing sets given by

$$\begin{array}{ccc}
 1 & \xrightarrow{(a, b)} & \{(x_0, \dots, x_n) : n \in \mathbb{N}, x_0 = a, x_n = b\} \\
 \downarrow & & \downarrow \text{apply } f_0 \\
 1 & \xrightarrow{(fa, fb)} & \{(y_0, \dots, y_n) : n \in \mathbb{N}, y_0 = fa, y_n = fb\}
 \end{array}$$

and the components are naturality squares for u . Thus by lemma(32) η is cartesian. Note that using the distributivity of E one has a canonical isomorphism

$$(\Gamma E)^2(X)(a, b) \cong \coprod_{(x_{ij})_{ij}} \prod_i E_i \prod_j E_j X(x_{ij-1}, x_{ij})$$

where the coproduct is taken over the set of composable doubly-indexed sequences starting at a and finishing at b . Unpacking in these terms one can see that in the case of μ 's hom naturality square, the underlying square of summand indexing sets is

$$\begin{array}{ccc}
 \{(x_{ij})_{ij} : x_0 = a, x_n = b\} & \xrightarrow{\text{concatenate}} & \{(x_0, \dots, x_n) : n \in \mathbb{N}, x_0 = a, x_n = b\} \\
 \text{apply } f_0 \downarrow & & \downarrow \text{apply } f_0 \\
 \{(y_{ij})_{ij} : y_0 = fa, y_n = fb\} & \xrightarrow{\text{concatenate}} & \{(y_0, \dots, y_n) : n \in \mathbb{N}, y_0 = fa, y_n = fb\}
 \end{array}$$

in which concatenation is that of *composable* sequences, that is, one identifies the last point of the i -th subsequence with the first point of the $(i+1)$ -th, which by definition of “composable doubly-indexed sequence” are equal as elements of X_0 or Y_0 . This square is easily seen to be a pullback. The components of μ 's hom naturality square are naturality squares for σ . Thus by lemma(32) μ is cartesian.

Conversely suppose that (Γ, η, μ) is a cartesian monad. Then by the same argument as for the converse direction of (2), except with pullbacks in place of λ -filtered colimits, one may conclude that E preserves pullbacks. Note that for $X \in V$ the hom between 0 and 1 of the naturality component of $\eta_{(X)}$ is, modulo the canonical isomorphism $E_1 X \cong \Gamma E(X)(0, 1)$, just $E_1 X$, and so u 's cartesianness follows from that of η by remark(21). Suppose that $(X_1, \dots, X_n)$ is a sequence of objects of V . Denote by $\text{sd}(X_i)_i$ the set of subdivisions of $(X_i)_i$ into a sequence of sequences. A typical element is a sequence of sequences (X_{ij}) where $1 \leq i \leq k$, $1 \leq j \leq n_i$ and $n_1 + \dots + n_k = n$, such that sequence obtained by concatenation is

$(X_1, \dots, X_n)$. Then modulo the canonical isomorphism

$$(\Gamma E)^2(X_1, \dots, X_n) \cong \coprod_{\text{sd}(X_i)_i} \mathbb{E}_i \mathbb{E}_j X_{ij}$$

the hom of the naturality component of $\mu_{(X_1, \dots, X_n)}$ between 0 and n is the map

$$(\sigma_{X_{ij}}) : \coprod_{\text{sd}(X_i)_i} \mathbb{E}_i \mathbb{E}_j X_{ij} \rightarrow \mathbb{E}_i X_i$$

and thus by remark(21), these maps are cartesian natural in the X_i . By lemma(2) V is extensive, and so the $\sigma_{X_{ij}}$ are cartesian natural in the X_{ij} as required. $\square$

4.6. Local right adjointness. We now proceed to the task of proving that the construction Γ is compatible with l.r.a-ness. For this we first require two lemmas. We assume familiarity with the notion of “generic morphism” and the alternative formulation of l.r.a-ness in terms of generics as described in [26] proposition(2.6).

Lemma 33. *Let $R : V \rightarrow W$ be a functor, V be cocomplete, U be a small dense full subcategory of W , and $L : U \rightarrow V$ be a partial left adjoint to R , that is to say, one has isomorphisms $W(S, RX) \cong V(LS, X)$ natural in $S \in U$ and $X \in V$. Defining $\overline{L} : W \rightarrow V$ as the left kan extension of L along the inclusion $I : U \rightarrow W$, one has $\overline{L} \dashv R$.*

PROOF. Denoting by $p : I/Y \rightarrow U$ the canonical forgetful functor for $Y \in W$ and recalling that $\overline{L}Y = \text{colim}(Lp)$, one obtains the desired natural isomorphism as follows

$$\begin{aligned} V(\overline{L}Y, X) &\cong [I/Y, V](Lp, \text{const}(X)) \cong \lim_{f \in I/Y} V(L(\text{dom}(f)), X) \\ &\cong \lim_f W(\text{dom}(f), RX) \cong \mathcal{B}(Y, RX) \end{aligned}$$

for all $X \in V$. $\square$

Lemma 34. *Let $T : V \rightarrow W$ be a functor, V be cocomplete and W have a small dense subcategory U . Then T is a local right adjoint iff every $f : S \rightarrow TX$ with $S \in U$ admits a generic factorisation. If in addition V has a terminal object denoted 1 , then generic factorisations in the case $X = 1$ suffice.*

PROOF. For the first statement ($\Rightarrow$) is true by definition so it suffices to prove the converse. The given generic factorisations provide a partial left adjoint $L : I/TX \rightarrow V$ to $T_X : V/X \rightarrow W/TX$ where I is the inclusion of U . Now I/TX is a small dense subcategory of W/TX , and so by the previous lemma L extends to a genuine left adjoint to T_X . In the case where V has 1 one requires only generic factorisations in the case $X = 1$ by the results of [26] section(2). $\square$

The analogous result for presheaf categories, with the representables forming the chosen small dense subcategory, was discussed in [26] section(2). With these results in hand we may now exhibit the

PROOF. (of theorem(29)(4))

The aspects of this result involving the cartesianness of the units, multiplication and substitution are covered already by (3). Suppose that E is l.r.a. Let $\mathcal{D}$ be a small dense subcategory of V consisting of λ -presentable connected objects. By lemma(34) and lemma(24) it suffices to exhibit generic factorisations of maps

$$f : S \rightarrow \Gamma E 1$$

where S is either 0 or (D) for some $D \in \mathcal{D}$. In the case where S is 0 the first arrow in the composite

$$0 \longrightarrow \Gamma E 0 \xrightarrow{\Gamma E t} \Gamma E 1$$

is generic because 0 is the initial V -graph with one object (and t here is the unique map). In the case where $S = (D)$, to give f is to give a map $f' : D \rightarrow E_n 1$ in V since D is connected. Since E is a local right adjoint, E_n is too and so one can generically factor f' to obtain

$$D \xrightarrow{g'_f} E_i Z_i \xrightarrow{E_i t} E_n 1$$

from which we obtain the generic factorisation

$$(D) \xrightarrow{g_f} \Gamma E Z \xrightarrow{\Gamma E t} \Gamma E 1$$

where $Z = (Z_1, \dots, Z_n)$, the object map of g_f is given by $0 \mapsto 0$ and $1 \mapsto n$, and the hom map of g_f is g'_f composed with the coproduct inclusion.

Conversely suppose that ΓE is l.r.a. It suffices by lemma(34) to exhibit a generic factorisation for maps of the form on the left in

$$f : Y \rightarrow E(X_1, \dots, X_n) \quad f' : (Y) \rightarrow \Gamma E(X_1, \dots, X_n)$$

where Y is connected. Such an f determines f' as in the previous display unique with object map $(0, 1) \mapsto (0, n)$ and hom map between 0 and 1 given by f , modulo the canonical isomorphism $E(X_1, \dots, X_n) \cong \Gamma E(X_1, \dots, X_n)(0, n)$ that we described already in the proof of (2).

Consider a factorisation

$$(Y) \xrightarrow{g} \Gamma E Z \xrightarrow{\Gamma E h} \Gamma E(X_1, \dots, X_n)$$

of f' . The object map of h partitions the objects of Z into $n+1$ subsets $Z_{(0)}, \dots, Z_{(n)}$. The strict initiality of $\emptyset$ and the definition of $(X_1, \dots, X_n) \in \mathcal{G}V$ ensures that the only homs of Z that are possibly non-initial, are those between a and b living in consecutive cells of this partition. Thus in addition to this partition h amounts to maps $h_{a,b} : Z(a, b) \rightarrow X_i$ for all $a \in Z_{(i-1)}$ and $b \in Z_{(i)}$. The connectedness of Y ensures that the hom map of g between 0 and 1 factors through a unique summand of the appropriate hom of $\Gamma E Z$. Thus the data of g comes down to: $1 \leq i \leq j \leq n$, $c_r \in Z_{(r)}$ for $i \leq r \leq j$ and a map $g_{0,1} : Y \rightarrow \bigcup_{i < r \leq j} Z(c_{r-1}, c_r)$. Consider the canonical inclusion

$$\bar{c} : (Z(c_i, c_{i+1}), \dots, Z(c_{j-1}, c_j)) \hookrightarrow Z$$

and note that by the above description one may factor g as

$$(Y) \xrightarrow{g'} \Gamma E(Z(c_i, c_{i+1}), \dots, Z(c_{j-1}, c_j)) \xrightarrow{\Gamma E \bar{c}} \Gamma E Z.$$

If g were in fact generic then $\bar{c}$ would have a section and thus be an isomorphism. It follows that any generic factorisation of f' is necessarily of the form

$$(Y) \xrightarrow{g} \Gamma E(X'_1, \dots, X'_n) \xrightarrow{\Gamma E(h_1, \dots, h_n)} \Gamma E(X_1, \dots, X_n)$$

for $h_i : X'_i \rightarrow X_i$ in V . Moreover it is easily shown that the hom of this factorisation between 0 and 1 gives a generic factorisation for the original map f , thereby exhibiting E as l.r.a. $\square$

5. The 2-functoriality of the multitensor to monad construction

5.1. The monad to multitensor construction. For a category V a monad (T, η, μ) over **Set** on $\mathcal{G}V$ is a monad on

$$(-)_0 : \mathcal{G}V \rightarrow \mathbf{Set}$$

in the 2-category **CAT/Set**. That is, the functor T doesn't affect the object sets and similarly for maps, and moreover the components of η and μ are identities on objects. Recall from section(3.1) that if V has an initial object then one can regard any sequence of objects $(Z_1, \dots, Z_n)$ of V as a V -graph. This is clearly functorial in the Z_i . Moreover for $1 \leq a \leq b \leq n$ one has subsequence inclusions

$$(Z_a, \dots, Z_b) \hookrightarrow (Z_1, \dots, Z_n)$$

defined in the obvious way, with the object map preserving successor and $0 \mapsto (a - 1)$, and the hom maps being identities. This enables us to construct a multitensor on V from T , essentially by applying T to sequences and looking at the homs.

Explicitly one defines this associated multitensor $(\bar{T}, \bar{\eta}, \bar{\mu})$ as follows. The n -ary tensor product is defined by

$$\bar{T}(Z_1, \dots, Z_n) := T(Z_1, \dots, Z_n)(0, n).$$

Recall that for $Z \in V$, (Z) is the V -graph with object set $\{0, 1\}$, hom between 0 and 1 equal to Z , and other homs initial. The unit $\bar{\eta}_Z : Z \rightarrow \bar{T}_1 Z$ is the hom map of $\eta_{(Z)}$ between 0 and 1. In order to define the substitution, note that given objects Z_{ij} of V where $1 \leq i \leq k$ and $1 \leq j \leq n_i$, one has a map

$$\tilde{\tau}_{Z_{ij}} : (\bigvee_{1 \leq j \leq n_1} \bar{T} Z_{1j}, \dots, \bigvee_{1 \leq j \leq n_k} \bar{T} Z_{kj}) \rightarrow T(Z_{11}, \dots, Z_{kn_k})$$

given on objects by $0 \mapsto 0$ and $i \mapsto (i, n_i)$ for $1 \leq i \leq k$. The hom map between $(i - 1)$ and i is the hom map of

$$Ts_i : T(Z_{i1}, \dots, Z_{in_i}) \rightarrow T(Z_{11}, \dots, Z_{kn_k})$$

between 0 and n_i , where s_i is the i -th subsequence inclusion. The component $\bar{\mu}_{Z_{ij}}$ is defined to be the hom map of $\mu \circ T(\tilde{\tau}_{Z_{ij}})$ between 0 and k .

In order to understand why $(\bar{T}, \bar{\eta}, \bar{\mu})$ form a multitensor, it is worthwhile to take a more conceptual approach. This begins with the observation that a sequence $(Z_1, \dots, Z_n)$ of objects of V may be viewed as a cospan

$$0 \xrightarrow{b} (X_1, \dots, X_n) \xleftarrow{t} 0$$

in $\mathcal{G}V$ in which b picks out the “bottom” object 0 and t picks out the “top” object n . Moreover pushout composition

$$\begin{array}{ccccc} & & 0 & & \\ & \swarrow b & & \searrow b & \\ 0 & & (Y_1, \dots, Y_m) & \xrightarrow{po} & (Z_1, \dots, Z_n) \\ & \nwarrow t & & \swarrow t & \\ & & 0 & & 0 \end{array}$$

$$(Y_1, \dots, Y_m, Z_1, \dots, Z_n)$$

of such cospans in $\mathcal{G}V$ corresponds, as shown, to concatenation of sequences. These pushouts are special in that they only require an initial object in V for their construction.

Pushout composition in $\mathcal{GV}$ of general cospans of the form

$$0 \longrightarrow X \longleftarrow 0$$

require coproducts in V for their construction. Note that such cospans are, as already pointed out in section(3.1), nothing more than bipointed V -graphs. Thus when V has coproducts, pushout composition of cospans endows the category $\mathcal{G}(t_V)_{\bullet}^{\times 2}$ of bipointed V -graphs with a monoidal structure whose tensor product we denote as “ $*$ ”. Moreover given a monad (T, η, μ) on $\mathcal{GV}$ over **Set**, one obtains a monoidal monad $(T_{\bullet}, \eta_{\bullet}, \mu_{\bullet})$ on $\mathcal{G}(t_V)_{\bullet}^{\times 2}$. The underlying endofunctor

$$T_{\bullet} : \mathcal{G}(t_V)_{\bullet}^{\times 2} \rightarrow \mathcal{G}(t_V)_{\bullet}^{\times 2} \quad (X, a, b) \mapsto (TX, a, b)$$

as object map as described in the previous display. In terms of cospans, this is just the application of T to cospans plus composition with the unique identity-on-objects $0 \rightarrow T0$ in order to get an endocospans of 0. The monoidal functor coherences for $T_{\bullet}$ are the maps that give the obstruction to T preserving the pushouts involved. The data $(\eta_{\bullet}, \mu_{\bullet})$ are defined in the evident way from (η, μ) .

The assignation of a cospan/bipointed V -graph from a sequence may done in two steps

$$(Z_1, \dots, Z_n) \in MV \mapsto (((Z_1), 0, 1), \dots, ((Z_n), 0, 1)) \mapsto ((Z_1, \dots, Z_n), 0, n)$$

and so is the object map of the composite

$$MV \xrightarrow{ML_V} M\mathcal{G}(t_V)_{\bullet}^{\times 2} \xrightarrow{*} \mathcal{G}(t_V)_{\bullet}^{\times 2}.$$

Thus one can view the functor $\overline{T} : MV \rightarrow V$ in more conceptual terms as the composite

$$MV \xrightarrow{ML_V} M\mathcal{G}(t_V)_{\bullet}^{\times 2} \xrightarrow{*} \mathcal{G}(t_V)_{\bullet}^{\times 2} \xrightarrow{T_{\bullet}} \mathcal{G}(t_V)_{\bullet}^{\times 2} \xrightarrow{\varepsilon_V} V.$$

Observe that $(T_{\bullet}, \eta_{\bullet}, \mu_{\bullet})$ is a monoidal monad and $L_V \dashv \varepsilon_V$. Moreover in general one has

Lemma 35. (1) *Let (E, u, σ) be a multitensor on V and (T, η, μ) be a monoidal monad on (V, E) with monoidal functor coherences for T written as*

$$\tau_{X_i} : E_i T X_i \rightarrow T E_i X_i.$$

Then (F, u', σ') defines another multitensor on V where $F_i X_i = T E_i X_i$, $u'_X = \eta_{E_1 X} u_X$ and σ' is the composite

$$T E_i T E_j \xrightarrow{T \tau E} T^2 E_i E_j \xrightarrow{\mu \sigma} T E_{ij}.$$

(2) *Let (E, u, σ) be a multitensor on V and $L \dashv R : V \rightarrow W$ with unit η and counit ε . Then (F, u', σ') defines multitensor on W where $F_i X_i = R E_i L X_i$, $u' = (R u M) \eta$ and σ' is the composite*

$$R E_i L R E_j L \xrightarrow{R E_i \varepsilon_j L} R E_i E_j L \xrightarrow{R \sigma L} R E_{ij} L$$

whose proof is an easy exercise in the definitions involved. Starting with the monoidal structure $*$ on $\mathcal{G}(t_V)_{\bullet}^{\times 2}$, apply (1) to obtain the multitensor $T_{\bullet}*$ on $\mathcal{G}(t_V)_{\bullet}^{\times 2}$, and then apply (2) to this using the adjunction $L_V \dashv \varepsilon_V$. It is straight forward to verify directly that the unit and substitution of the resulting multitensor coincides with $(\overline{\eta}, \overline{\mu})$ as defined above. Thus we have

Proposition 36. *Let V be a category with coproducts and (T, η, μ) be a monad on $\mathcal{G}V$ over **Set**. Then $(\overline{T}, \overline{\eta}, \overline{\mu})$ defines a multitensor on V .*

Remark 37. Note that the multitensor $(\overline{T}, \overline{\eta}, \overline{\mu})$ played an implicit role the proofs of the converse implications of theorem(29)(2)-(4). The reason for this is that one has a canonical isomorphism $E \cong \overline{\Gamma E}$ of multitensors. The isomorphism at the level of functors $MV \rightarrow V$ was described in the proof of theorem(29)(2), and the reader will easily verify the compatibility of this isomorphism with the unit and substitution maps.

5.2. Characterisation of monads coming from multitensors. In this section we characterise the monads of the form $(\mathcal{G}V, \Gamma E)$ as those monads on $\mathcal{G}V$ over **Set** which are distributive and path-like in the sense to be defined below. Throughout this section V is a category with coproducts.

Given a V -graph X and sequence $x = (x_0, \dots, x_n)$ of objects of X , one can define the morphism

$$\overline{x} : (X(x_0, x_1), X(x_1, x_2), \dots, X(x_{n-1}, x_n)) \rightarrow X$$

whose object map is $i \mapsto x_i$, and whose hom map between $(i-1)$ and i is the identity. For all such sequences x one has

$$T(\overline{x})_{0,n} : \overline{T}_i X(x_{i-1}, x_i) \rightarrow TX(x_0, x_n)$$

and so taking all sequences x starting at a and finishing at b one induces the canonical map

$$\pi_{X,a,b} : \coprod_{a=x_0, \dots, x_n=b} \overline{T}_i X(x_{i-1}, x_i) \rightarrow TX(a, b).$$

Definition 38. Let V be a category with coproducts and (T, η, μ) be a monad on $\mathcal{G}V$ over **Set**. Then T is said to be *path-like* when for all $X \in \mathcal{G}V$ and $a, b \in X_0$, the maps $\pi_{X,a,b}$ are isomorphisms.

Example 39. Let $V = \mathbf{Set}$ and T be the free category endofunctor of **Gph**. For any graph X and $a, b \in X_0$, the hom $TX(a, b)$ is by definition the set of paths in X from a to b . Each path determines a sequence $x = (x_0, \dots, x_n)$ of objects of X such that $x_0 = a$ and $x_n = b$, by reading off the objects of X as they are visited by the given path. Conversely for a sequence $x = (x_0, \dots, x_n)$ of objects of X such that $x_0 = a$ and $x_n = b$, $T\overline{x}$ identifies the elements $\overline{T}_i X(x_{i-1}, x_i)$ with those paths in X from a to b whose associated sequence is x . Thus T is path-like.

Proposition 40. *Let V have small coproducts and (T, η, μ) be a path-like monad on $\mathcal{G}V$ over **Set**. Then $\mathcal{G}(V)^T \cong \overline{T}\text{-Cat}$.*

PROOF. Let X be a V -graph. To give an identity on objects map $a : TX \rightarrow X$ is to give maps $a_{y,z} : TX(y, z) \rightarrow X(y, z)$. By path-likeness these amount to giving for each $n \in \mathbb{N}$ and $x = (x_0, \dots, x_n)$ such that $x_0 = y$ and $x_n = z$, a map

$$a_x : \overline{T}_i X(x_{i-1}, x_i) \rightarrow X(y, z)$$

since $\overline{T}_i X(x_{i-1}, x_i) = Tx^*X(0, n)$, that is $a_x = a_{y,z}T\overline{x}_{0,n}$. When $n = 1$, for a given $y, z \in X_0$, x can only be the sequence (y, z) . The naturality square for η at $\overline{x}$ implies that $\{\eta_X\}_{y,z} = T\overline{x}_{0,1}\{\eta_{(X(y,z))}\}_{0,1}$, and the definition of $(\overline{})$ says that $\{\eta_{(X(y,z))}\}_{0,1} = \overline{\eta}_{X(y,z)}$. Thus to say that a map $a : TX \rightarrow X$ satisfies the unit law of a T -algebra is to say that a is the identity on objects and that the a_x described above satisfy the unit axioms of a $\overline{T}$ -category.

To say that a satisfies the associative law is to say that for all $y, z \in X_0$,

$$(6) \quad \begin{array}{ccc} T^2X(y, z) & \xrightarrow{\{\mu_X\}_{y,z}} & TX(y, z) \\ Tx_{y,z} \downarrow & & \downarrow a_{y,z} \\ TX(y, z) & \xrightarrow{a_{y,z}} & X(y, z) \end{array}$$

commutes. Given $x = (x_0, \dots, x_n)$ from X with $x_0 = y$ and $x_n = z$, and $w = (w_0, \dots, w_k)$ from $T(X(x_0, x_1), \dots, X(x_{n-1}, x_n))$ with $w_0 = 0$ and $w_k = n$, one can consider the composite map $T(\overline{x})_{0,n}T(\overline{w})_{0,k}$, and since the coproduct of coproducts is a coproduct, all such maps for x and w such that $x_0 = y$ and $x_n = z$ form a coproduct cocone. Precomposing (6) with the coproduct inclusions gives the commutativity of

$$\begin{array}{ccc} \overline{T}_i \overline{T}_j X(x_{ij-1}, x_{ij}) & \xrightarrow{\overline{\mu}} & \overline{T}_{ij} X(x_{ij-1}, x_{ij}) \\ \overline{T}_i a \downarrow & & \downarrow a_x \\ \overline{T}_i X(x_{w_{i-1}}, x_{w_i}) & \xrightarrow{a_w} & X(y, z) \end{array}$$

and conversely by the previous sentence if these squares commute for all x and w , then one recovers the commutativity of (6). This completes the description of the object part of $\mathcal{G}(V)^T \cong \overline{T}\text{-Cat}$.

Let (X, a) and (X', a') be T -algebras and $F : X \rightarrow X'$ be a V -graph morphism. To say that F is a T -algebra map is a condition on the maps $F_{y,z} : X(y, z) \rightarrow X'(Fy, Fz)$ for all $y, z \in X_0$, and one uses path-likeness in the obvious way to see that this is equivalent to saying that the $F_{y,z}$ are the hom maps for a $\overline{T}$ -functor. $\square$

Definition 41. Let V be a category with coproducts and (T, η, μ) be a monad on $\mathcal{G}V$ over **Set**. Then T is said to be *distributive* when $\overline{T}$ is a distributive multitensor.

Theorem 42. Let V have coproducts. Then a monad T on $\mathcal{G}V$ over **Set** is of the form $(\mathcal{G}V, \Gamma E)$ iff it is distributive and path-like, and in this case E is recovered as $\overline{T}$.

PROOF. Suppose that T is distributive and path-like. Since $\overline{T}$ is distributive the morphisms $\pi_{X,a,b}$ are the hom maps of the components of a natural transformation $\pi : \Gamma \overline{T} \rightarrow T$, which is easily seen to be compatible with the monad structures. Since T is path-like, this is an isomorphism. The converse follows from remark(37). $\square$

5.3. 2-functoriality of Γ . As the lax-algebras of a 2-monad M lax monoidal categories form a 2-category $\text{Lax-}M\text{-Alg}$. See [19] for a complete description of the 2-category of lax algebras for an arbitrary 2-monad. Explicitly a lax monoidal

functor between lax monoidal categories (V, E) and (W, F) consists of a functor $H : V \rightarrow W$, and maps

$$\psi_{X_i} : F_i H X_i \rightarrow H E_i X_i$$

natural in the X_i such that

$$\begin{array}{ccc} H X & \xrightarrow{u_{HX}} & F_1 H X \\ H u_X \searrow & & \swarrow \psi_X \\ & H E_1 X & \end{array} \quad \begin{array}{ccccc} F_i F_j H X_{ij} & \xrightarrow{F_i \psi} & F_i H E_j X_{ij} & \xrightarrow{\psi E_j} & H E_i E_j X_{ij} \\ \sigma H \searrow & & \swarrow H \sigma & & \\ & F_{ij} H X_{ij} & \xrightarrow{\psi} & H E_{ij} X_{ij} & \end{array}$$

commute for all X and X_{ij} in V . A monoidal natural transformation between lax monoidal functors

$$(H, \psi), (K, \kappa) : (V, E) \rightarrow (W, F)$$

consists of a natural transformation $\phi : H \rightarrow K$ such that

$$\begin{array}{ccc} F_i H X_i & \xrightarrow{\psi} & H E_i X_i \\ F_i \phi \downarrow & & \downarrow \phi E_i \\ F_i K X_i & \xrightarrow{\kappa} & K E_i X_i \end{array}$$

commutes for all X_i . We denote by **DISTMULT** the 2-category **DISTMULT** of distributive multitensors. It is the full sub-2-category of **Lax- M -Alg** consisting of the (V, E) such that V has coproducts and E is distributive.

For any 2-category $\mathcal{K}$ recall the 2-category $\mathbf{MND}(\mathcal{K})$ from [24] of monads in $\mathcal{K}$. Another way to describe this very canonical object is that it is the 2-category of lax algebras of the identity monad on $\mathcal{K}$. Explicitly the 2-category $\mathbf{MND}(\mathbf{CAT})$ has as objects pairs (V, T) where V is a category and T is a monad on V . An arrow $(V, T) \rightarrow (W, S)$ is a pair consisting of a functor $H : V \rightarrow W$ and a natural transformation $\psi : SH \rightarrow HT$ satisfying the obvious 2 axioms: these are just the “unary” analogues of the axioms for a lax monoidal functor written out above. For example, any lax monoidal functor (H, ψ) as above determines a monad functor $(H, \psi_1) : (V, E_1) \rightarrow (W, F_1)$. A monad transformation between monad functors

$$(H, \psi), (K, \kappa) : (V, T) \rightarrow (W, S)$$

consists of a natural transformation $\phi : H \rightarrow K$ satisfying the obvious axiom. For example a monoidal natural transformation ϕ as above is a monad transformation $(H, \psi_1) \rightarrow (K, \kappa_1)$.

In fact as we are interested in monads over **Set**, we shall work not with $\mathbf{MND}(\mathbf{CAT})$ but rather with $\mathbf{MND}(\mathbf{CAT}/\mathbf{Set})$. An arrow $(V, T) \rightarrow (W, S)$ of this 2-category is a pair (H, ψ) as in the case of $\mathbf{MND}(\mathbf{CAT})$, with the added condition that ψ 's components are the identities on objects, and similarly the 2-cells of $\mathbf{MND}(\mathbf{CAT}/\mathbf{Set})$ come with an extra identity-on-object condition.

We shall now exhibit the 2-functor

$$\Gamma : \mathbf{DISTMULT} \rightarrow \mathbf{MND}(\mathbf{CAT}/\mathbf{Set})$$

which on objects is given by $(V, E) \mapsto (GV, \Gamma E)$. Let $(H, \psi) : (V, E) \rightarrow (W, F)$ be a lax monoidal functor between distributive lax monoidal categories. Then for

$X \in \mathcal{G}V$ and $a, b \in X_0$, we define the hom map $\Gamma(\psi)_{X,a,b}$ as

$$\begin{array}{ccc}
 \coprod_{a=x_0, \dots, x_n=b} F H X(x_{i-1}, x_i) & \xrightarrow{\coprod \psi} & \coprod_{a=x_0, \dots, x_n=b} H E X(x_{i-1}, x_i) \\
 \searrow \Gamma(\psi)_{X,a,b} & = & \swarrow \text{obst.} \\
 H \coprod_{a=x_0, \dots, x_n=b} E X(x_{i-1}, x_i) & &
 \end{array}$$

where “obst.” denotes the obstruction map to H preserving coproducts. It follows easily from the definitions that $(\mathcal{G}H, \Gamma(\psi))$ as defined here satisfies the axioms of a monad functor. Moreover given a monoidal natural transformation $\phi : (H, \psi) \rightarrow (K, \kappa)$, it also follows easily from the definitions that

$$\mathcal{G}\phi : (\mathcal{G}H, \Gamma(\psi)) \rightarrow (\mathcal{G}K, \Gamma(\kappa))$$

is a monad transformation. It is also straight-forward to verify that these assignments are 2-functorial.

Lax algebras of a 2-monad organise naturally into *two* different 2-categories depending on whether one takes lax or oplax algebra morphisms. So in particular one has the 2-category $\text{OpLax-}M\text{-Alg}$ of lax monoidal categories, *oplax*-monoidal functors between them and monoidal natural transformations between those. The coherence ψ for an oplax $(H, \psi) : (V, E) \rightarrow (W, F)$ goes in the other direction, and so its components look like this:

$$\psi_{X_i} : H E X_i \rightarrow F H X_i.$$

The reader should easily be able to write down explicitly the two coherence axioms that this data must satisfy, as well as the condition that must be satisfied by a monoidal natural transformation between oplax monoidal functors. Similarly there is a dual version $\text{OpMND}(\mathcal{K})$ of the 2-category $\text{MND}(\mathcal{K})$ of monads in a given 2-category $\mathcal{K}$ discussed above [24]. An arrow $(V, T) \rightarrow (W, S)$ of $\text{OpMND}(\mathbf{CAT})$ consists of a functor $H : V \rightarrow W$ and a natural transformation $\psi : HT \rightarrow SH$ satisfying the two obvious axioms. An arrow of $\text{OpMND}(\mathbf{CAT})$ is called a monad opfunctor. As before $\text{OpMND}(\mathbf{CAT}/\mathbf{Set})$ differs from $\text{MND}(\mathbf{CAT}/\mathbf{Set})$ in that all the categories involved come with a functor into $\mathbf{Set}$, and all the functors and natural transformations involved are compatible with these forgetful functors.

When defining the one-cell map of Γ above we were helped by the fact that the coproduct preservation obstruction went the right way: see the definition of the monad functor $(\mathcal{G}H, \Gamma\psi)$ above. This time however we will not be so lucky. For this reason we define the 2-category OpDISTMULT to be the locally full sub-2-category of $\text{OpLax-}M\text{-Alg}$ consisting of the distributive lax monoidal categories, and the oplax monoidal functors (H, ψ) such that H preserves coproducts. Thus we can define

$$\Gamma' : \text{OpDISTMULT} \rightarrow \text{OpMND}(\mathbf{CAT}/\mathbf{Set})$$

on objects by $(V, E) \mapsto (\mathcal{G}V, \Gamma E)$. Let $(H, \psi) : (V, E) \rightarrow (W, F)$ be an oplax monoidal functor between distributive lax monoidal categories. Then for $X \in \mathcal{G}V$ and $a, b \in$

X_0 , we define the hom map $\Gamma'(\psi)_{X,a,b}$ as

$$\begin{array}{ccc} H \coprod_{a=x_0, \dots, x_n=b} \coprod_i E X(x_{i-1}, x_i) & \xrightarrow{\cong} & \coprod_{a=x_0, \dots, x_n=b} H \coprod_i E X(x_{i-1}, x_i) \\ & \searrow \Gamma'(\psi)_{X,a,b} \quad \quad \quad \swarrow \coprod \psi & \\ & \coprod_{a=x_0, \dots, x_n=b} F H X(x_{i-1}, x_i) & \end{array}$$

It follows easily from the definitions that $(\mathcal{G}H, \Gamma'(\psi))$ as defined here satisfies the axioms of a monad opfunctor. Moreover given a monoidal natural transformation $\phi : (H, \psi) \rightarrow (K, \kappa)$, it also follows easily from the definitions that

$$\mathcal{G}\phi : (\mathcal{G}H, \Gamma'(\psi)) \rightarrow (\mathcal{G}K, \Gamma'(\kappa))$$

is a monad transformation. It is also straight-forward to verify that these assignments are 2-functorial.

5.4. Properties of the 2-functor Γ .

Proposition 43. *Γ and Γ' are locally fully faithful 2-functors.*

PROOF. We will verify that Γ is locally fully faithful; the proof for Γ' is similar. Let $(H, \psi), (K, \kappa) : (V, E) \rightarrow (W, F)$ be lax monoidal functors between distributive lax monoidal categories. Given $\phi : \mathcal{G}H \rightarrow \mathcal{G}K$ so that $\phi : (\mathcal{G}H, \Gamma(\psi)) \rightarrow (\mathcal{G}K, \Gamma(\kappa))$ is a monad 2-cell, we must exhibit a unique monoidal natural transformation $\phi' : H \rightarrow K$ such that $\mathcal{G}\phi' = \phi$. By proposition(17) there is a unique $\phi' : H \rightarrow K$ such that $\mathcal{G}\phi' = \phi$, and from the proof of proposition(17) this is defined by $\phi'_Z = \phi_{(Z),0,1}$. So it suffices to show that ϕ satisfies the monad 2-cell axiom iff ϕ' satisfies the monoidal naturality axiom. The monad 2-cell axiom says the outside of

$$\begin{array}{ccccc} \coprod_i F H X(x_{i-1}, x_i) & \xrightarrow{\coprod \psi} & \coprod_i H \coprod E X(x_{i-1}, x_i) & \xrightarrow{\text{obstn}} & H \coprod_i E X(x_{i-1}, x_i) \\ \downarrow \coprod F \phi' & & \downarrow \coprod \phi' & & \downarrow \phi_{\Gamma E(X),a,b} \\ \coprod_i F H X(x_{i-1}, x_i) & \xrightarrow{\coprod \kappa} & \coprod_i K \coprod E X(x_{i-1}, x_i) & \xrightarrow{\text{obstn}} & K \coprod_i E X(x_{i-1}, x_i) \end{array}$$

commutes for all $X \in \mathcal{G}V$ and $a, b \in X_0$, and where all the coproducts are taken over all sequences $a = x_0, \dots, x_n = b$. Since $\phi_{\Gamma E(X),a,b} = \phi'_{\Gamma E(X)(a,b)}$, the right hand square commutes by the naturality of the obstruction maps. Monoidal naturality of ϕ' says that for all $(Z_1, \dots, Z_n)$

$$\begin{array}{ccc} F H Z_i & \xrightarrow{\psi} & H \coprod E Z_i \\ \downarrow F \phi' & & \downarrow \phi' \\ F K Z_i & \xrightarrow{\psi'} & K \coprod E Z_i \end{array}$$

commutes, which implies that the left hand square above commutes, and so monoidal naturality of ϕ' implies the monad 2-cell axiom for ϕ . For the converse take $X = (Z_1, \dots, Z_n)$, $a = 0$ and $b = n$. In this case the coproduct involved in the monad 2-cell axiom has only one non-trivial summand, that for the sequence $(0, 1, \dots, n)$. Thus the obstruction maps are isomorphisms, and the left hand square is exactly the monoidal naturality axiom for ϕ' . $\square$

While Γ and Γ' aren't themselves 2-fully faithful, proposition(44) is a useful related statement which is true. By definition Γ and Γ' fit into commutative squares

$$\begin{array}{ccc} \text{DISTMULT} & \xrightarrow{\Gamma} & \text{MND}(\mathbf{CAT}/\mathbf{Set}) \\ \downarrow & & \downarrow \\ \mathbf{CAT} & \xrightarrow{\mathcal{G}_1} & \mathbf{CAT}/\mathbf{Set} \end{array} \quad \begin{array}{ccc} \text{OpDISTMULT} & \xrightarrow{\Gamma'} & \text{OpMND}(\mathbf{CAT}/\mathbf{Set}) \\ \downarrow & & \downarrow \\ \mathbf{CAT} & \xrightarrow{\mathcal{G}_1} & \mathbf{CAT}/\mathbf{Set} \end{array}$$

in which the vertical arrows are the obvious forgetful 2-functors. Let us write $\mathcal{G}\text{-MND}$ (resp. $\mathcal{G}\text{-OpMND}$) for the 2-categories obtained by pulling back $\mathcal{G}_1$ along the appropriate forgetful 2-functor, and by

$$\Psi : \text{DISTMULT} \rightarrow \mathcal{G}\text{-MND} \quad \Psi' : \text{OpDISTMULT} \rightarrow \mathcal{G}\text{-OpMND}$$

the induced 2-functors.

In more concrete terms an object of $\mathcal{G}\text{-MND}$ (or of $\mathcal{G}\text{-OpMND}$) is a pair (V, T) where V is a category with coproducts and T is a monad on $\mathcal{G}V$ over $\mathbf{Set}$. By definition and by theorem(42), we know that (V, T) is in the image of Ψ (or of Ψ') iff T is distributive and path-like. A morphism $(V, T) \rightarrow (W, S)$ in $\mathcal{G}\text{-MND}$ is a pair (H, ψ) where $H : V \rightarrow W$ is a functor, and ψ is 2-cell data making $(\mathcal{G}H, \psi) : (\mathcal{G}V, T) \rightarrow (\mathcal{G}W, S)$ a monad functor. Similarly, a morphism $(V, T) \rightarrow (W, S)$ in $\mathcal{G}\text{-OpMND}$ is a pair (H, ψ) where $H : V \rightarrow W$ is a coproduct preserving functor, and ψ is 2-cell data making $(\mathcal{G}H, \psi) : (\mathcal{G}V, T) \rightarrow (\mathcal{G}W, S)$ a monad opfunctor. A 2-cell $\phi : (H, \psi) \rightarrow (K, \kappa)$ of $\mathcal{G}\text{-MND}$ is a natural transformation $\phi : H \rightarrow K$, making $\mathcal{G}\phi : (\mathcal{G}H, \psi) \rightarrow (\mathcal{G}K, \kappa)$ a monad 2-cell, and the 2-cells of $\mathcal{G}\text{-OpMND}$ are described similarly.

Proposition 44. *Ψ and Ψ' are 2-fully faithful.*

PROOF. We shall prove that Ψ is 2-fully faithful; the proof for Ψ' is similar. By definition and proposition(43) Ψ is locally fully faithful. Thus it suffices to show that Ψ is fully faithful as a mere functor. This in turn amounts to showing that for any functor $H : V \rightarrow W$ between categories with coproducts, and any natural transformation $\psi : \Gamma(F)\mathcal{G}(H) \rightarrow \mathcal{G}(H)\Gamma(E)$ such that $(\mathcal{G}H, \psi) : (\mathcal{G}V, \Gamma E) \rightarrow (\mathcal{G}W, \Gamma F)$ is a monad functor, that there exists a unique $\psi' : FM(H) \rightarrow HE$ making $(H, \psi') : (V, E) \rightarrow (W, F)$ a lax monoidal functor such that $\Gamma\psi' = \psi$.

The homs of the components of ψ are maps in V of the form

$$\psi_{X,a,b} : \coprod_{a=x_0, \dots, x_n=b} F \coprod_i H X(x_{i-1}, x_i) \rightarrow H \coprod_{a=x_0, \dots, x_n=b} E X(x_{i-1}, x_i)$$

and in the case where $X = (Z_1, \dots, Z_n)$, $a = 0$ and $b = n$, the coproducts here have only one non-trivial summand, that for the sequence $(0, 1, \dots, n)$, and so we define

$$\psi'_{Z_i} := \psi_{(Z_1, \dots, Z_n), 0, n} : F \coprod_i H Z_i \rightarrow H \coprod_i E Z_i.$$

The lax monoidal functor coherence axioms for ψ' follow easily from the monad functor coherence axioms for ψ . To say that $\Gamma\psi' = \psi$ is to say that

$$\begin{array}{ccc} \coprod_{a=x_0, \dots, x_n=b} F_i H X(x_{i-1}, x_i) & \xrightarrow{\psi_{X,a,b}} & H \coprod_{a=x_0, \dots, x_n=b} E_i X(x_{i-1}, x_i) \\ & \searrow \coprod \psi' & \nearrow \text{obstn} \\ & \coprod_{a=x_0, \dots, x_n=b} H E_i X(x_{i-1}, x_i) & \end{array}$$

commutes for all $X \in \mathcal{G}V$ and $a, b \in X_0$, which is to say that

$$\begin{array}{ccc} F_i H X(x_{i-1}, x_i) & \xrightarrow{c_{x_i}} & \coprod_{a=x_0, \dots, x_n=b} F_i H X(x_{i-1}, x_i) \\ \psi' \downarrow & & \downarrow \psi_{X,a,b} \\ H E_i X(x_{i-1}, x_i) & \xrightarrow{H c_{x_i}} & H \coprod_{a=x_0, \dots, x_n=b} E_i X(x_{i-1}, x_i) \end{array}$$

commutes for all $X \in \mathcal{G}V$ and $a = x_0, \dots, x_n = b \in X_0$. But this last square is just the hom between a and b for the naturality square for ψ with respect to the canonical inclusion $(X(x_0, x_1), \dots, X(x_{n-1}, x_n)) \hookrightarrow X$. Finally we note that given ϕ making $(H, \phi) : (V, E) \rightarrow (W, F)$ a lax monoidal functor one has for $Z_1, \dots, Z_n \in V$

$$(\Gamma\phi)'_{Z_1, \dots, Z_n} = (\Gamma\phi)_{(Z_1, \dots, Z_n), 0, n} = \phi_{Z_i}$$

and so $\psi \mapsto \psi'$ is the inverse of the arrow map of Ψ . $\square$

5.5. Cartesian transformations. We now note that the above constructions are compatible with cartesian transformations.

Lemma 45. *Suppose that $H : V \rightarrow W$ is a pullback preserving functor between extensive categories. Then the obstruction maps*

$$\coprod_{i \in I} H X_i \rightarrow H \coprod_{i \in I} X_i$$

are cartesian-natural in the X_i .

PROOF. The naturality squares in question appear as the right hand square in

$$\begin{array}{ccccc} & & H c_{X_i} & & \\ & \curvearrowright & & \curvearrowright & \\ H X_i & \xrightarrow{c_{H X_i}} & \coprod H X_i & \xrightarrow{\text{obstn}} & H \coprod X_i \\ H f_i \downarrow & & \downarrow \coprod H f_i & & \downarrow H \coprod f_i \\ H Y_i & \xrightarrow{c_{H Y_i}} & \coprod H Y_i & \xrightarrow{\text{obstn}} & H \coprod Y_i \\ & \curvearrowright & & \curvearrowright & \\ & & H c_{Y_i} & & \end{array}$$

Since V is extensive and H preserves pullbacks it follows that outside square is a pullback for all $i \in I$. Since W is extensive it follows then that the right hand square is a pullback as required. $\square$

Proposition 46. (1) *Let $(H, \psi) : (V, E) \rightarrow (W, F)$ be a lax monoidal functor between distributive lax monoidal categories, and suppose that V and W*

are extensive and H preserves pullbacks. Then the following statements are equivalent:

- (a) ψ is a cartesian transformation.
 - (b) $\Gamma\psi$ is a cartesian transformation.
 - (c) $\Psi\psi$ is a cartesian transformation.
- (2) Let $(H, \psi) : (V, E) \rightarrow (W, F)$ be a coproduct preserving oplax monoidal functor between distributive lax monoidal categories, and suppose that W is extensive. Then the following statements are equivalent:
- (a) ψ is a cartesian transformation.
 - (b) $\Gamma'\psi$ is a cartesian transformation.
 - (c) $\Psi'\psi$ is a cartesian transformation.

PROOF. The statements that $\Gamma\psi$ cartesian iff $\Psi\psi$ is cartesian, and similarly for $\Gamma'\psi$ and $\Psi'\psi$, follows by definition. For (H, ψ) as in (1) note that by the extensivity of W the maps

$$\coprod_{a=x_0, \dots, x_n=b} \psi_{X(x_{i-1}, x_i)} : \coprod_{a=x_0, \dots, x_n=b} F_i H X(x_{i-1}, x_i) \rightarrow \coprod_{a=x_0, \dots, x_n=b} H E_i X(x_{i-1}, x_i)$$

are cartesian natural iff ψ is, and so (1) follows by lemma(45) and the definition of Γ . For (H, ψ) as in (2) one has by the extensivity of W that the cartesian naturality of

$$\coprod_{a=x_0, \dots, x_n=b} \psi_{X(x_{i-1}, x_i)} : \coprod_{a=x_0, \dots, x_n=b} H E_i X(x_{i-1}, x_i) \rightarrow \coprod_{a=x_0, \dots, x_n=b} F_i H X(x_{i-1}, x_i)$$

is equivalent to that of ψ , so that (2) follows from the definition of Γ' . $\square$

Recall that for a cartesian monad (T, η, μ) on a category V with pullbacks, a T -operad consists of another monad A on V together with a cartesian monad morphism $\alpha : A \rightarrow T$, that is to say, one has a natural transformation $\alpha : A \rightarrow T$ whose naturality squares are pullbacks, and is a morphism of monoids in the monoidal category $\text{End}(V)$. Given a cartesian monad T on $\mathcal{G}V$ over **Set**, a T -operad over **Set** is defined in the same way except that the natural transformation α lives over **Set**, which means that in addition α 's components are identities on objects. For instance for $T = \mathcal{T}_{\leq n}$ the strict n -category monad on $\mathcal{G}^n \mathbf{Set}$, T -operads over **Set** were called *normalised n -operads* of [3] and the terminology “normalised” was also used in [6]. Finally we note that Γ 's image is closed under cartesian monad maps.

Lemma 47. *Let V be a lextensive category and T be a cartesian monad on $\mathcal{G}V$ over **Set**. Let $\alpha : A \rightarrow T$ be a T -operad over **Set**.*

- (1) *If T is distributive then so is A .*
- (2) *If T is path-like then so is A .*

PROOF. (1): given an n -tuple $(X_1, \dots, X_n)$ of objects of V and a coproduct cocone

$$(c_j : X_{ij} \rightarrow X_i : j \in J)$$

where $1 \leq i \leq n$, we must show that the hom-maps

$$A(X_1, \dots, c_j, \dots, X_n)_{0,n} : A(X_1, \dots, X_{ij}, \dots, X_n)(0, n) \rightarrow A(X_1, \dots, X_i, \dots, X_n)(0, n)$$

form a coproduct cocone. For $j \in J$ we have a pullback square

$$\begin{array}{ccc} A(X_1, \dots, X_{ij}, \dots, X_n)(0, n) & \xrightarrow{A(X_1, \dots, c_j, \dots, X_n)_{0,n}} & A(X_1, \dots, X_i, \dots, X_n)(0, n) \\ \alpha \downarrow & \text{pb} & \downarrow \alpha \\ T(X_1, \dots, X_{ij}, \dots, X_n)(0, n) & \xrightarrow{T(X_1, \dots, c_j, \dots, X_n)_{0,n}} & T(X_1, \dots, X_i, \dots, X_n)(0, n) \end{array}$$

and by the distributivity of T the $T(X_1, \dots, c_j, \dots, X_n)_{0,n}$ form a coproduct cocone, and thus so do the $A(X_1, \dots, c_j, \dots, X_n)_{0,n}$ by the extensivity of V .

(2): given $X \in \mathcal{GV}$, $a, b \in X_0$ and a sequence $(x_0, \dots, x_n)$ of objects of X such that $x_0 = a$ and $x_n = b$, we have the map

$$A\overline{x}_{0,n} : A(X(x_0, x_1), \dots, X(x_{n-1}, x_n))(0, n) \rightarrow AX(a, b)$$

and we must show that these maps, where the x_i range over all sequences from a to b , form a coproduct cocone. By the path-likeness of T we know that the maps

$$T\overline{x}_{0,n} : T(X(x_0, x_1), \dots, X(x_{n-1}, x_n))(0, n) \rightarrow TX(a, b)$$

form a coproduct cocone, so we can use the cartesianness of α and the extensivity of V to conclude as in (1). $\square$

6. Distributive laws and monoidal monads

6.1. Distributing a monad over a multitensor. Let (V, E, u, σ) be a lax monoidal category. Recall that a *monoidal monad* on (V, E) is a monad on (V, E) in the 2-category of lax monoidal categories, lax monoidal functors and monoidal natural transformations. In more explicit terms this amounts to a monad (T, η, μ) on V together with coherence maps

$$\tau_{X_i} : E_i T X_i \rightarrow T E_i X_i$$

such that

$$\begin{array}{ccc} T & \xrightarrow{uT} & E_1 T \\ Tu \searrow & & \swarrow \tau \\ & T E_1 & \end{array} \qquad \begin{array}{ccccc} E_i E_j T & \xrightarrow{E_i \tau} & E_i T E_j & \xrightarrow{\tau E_j} & T E_i E_j \\ \sigma T \searrow & & \swarrow T \sigma & & \\ & E_{ij} T & \xrightarrow{\tau} & T E_{ij} & \end{array}$$

and

$$\begin{array}{ccc} E_1 & \xrightarrow{E_1 \eta} & E_1 T \\ \eta E_1 \searrow & & \swarrow \tau \\ & T E_1 & \end{array} \qquad \begin{array}{ccccc} E_i T^2 & \xrightarrow{\tau T} & T E_i T & \xrightarrow{T \tau} & T^2 E_i \\ E_i \mu \searrow & & \swarrow \mu E_i & & \\ & E_i T & \xrightarrow{\tau} & T E_i & \end{array}$$

commute. Ignoring the subscripts in the above data and axioms one can see immediately the formal resemblance with monad distributive laws. Restricting attention to singleton sequences of objects from V one has a monad distributive law of T over E_1 .

Given a multitensor E on a category V and a monad (S, η, μ) on $\mathcal{G}V$, a *lifting of S to E -Cat* is a monad (S', η', μ') on E -Cat such that $SU^E = U^E S'$, $\eta U^E = U^E \eta'$ and $\mu U^E = U^E \mu'$, where we recall that $U^E : E\text{-Cat} \rightarrow \mathcal{G}V$ is the forgetful functor.

The central results of the theory of monad distributive laws [7] provide an equivalence between monad distributive laws of a monad S over a monad T on some category V , with data giving an induced monad structure on the composite ST , with liftings of the monad S to a monad S' on the algebras of T . Moreover for any instance of such data one has an isomorphism $V^{ST} \cong (V^T)^{S'}$ commuting with the forgetful functors into V . The analogous results for monoidal monads are contained in

Theorem 48. *Let V be a category, (E, u, σ) be a multitensor on V and (T, η, μ) be a monad on V . Then there is a bijection between the following types of data:*

- (1) *Morphisms $\tau_{X_i} : E_i T X_i \rightarrow T E_i X_i$ of V providing the coherences making T into a monoidal monad.*
- (2) *Morphisms $\sigma'_{X_{ij}} : T E_i T E_j X_{ij} \rightarrow T E_{ij} X_{ij}$ of V providing the substitutions for a multitensor (TE, u', σ') where $u'_X = \eta_{E_1 X} u_X$, $\eta_E : E \rightarrow TE$ is a multitensor map, $Tu : T \rightarrow TE_1$ is a monad map, and the composite*

$$T E_i \xrightarrow{T u \eta E_i} T E_1 T E_i \xrightarrow{\sigma'} T E_i$$

is the identity.

if in addition V has coproducts and E is distributive, then the data (1)-(2) are also in bijection with

- (3) *Identity on object morphisms $\lambda_X : \Gamma(E)\mathcal{G}(T)(X) \rightarrow \mathcal{G}(T)\Gamma(E)(X)$ of $\mathcal{G}V$ providing a monad distributive law of $\mathcal{G}(T)$ over $\Gamma(E)$.*
- (4) *Liftings of the monad $\mathcal{G}(T)$ to a monad T' on E -Cat.*

and for any given instance of such data one has an isomorphism

$$(TE)\text{-Cat} \cong E\text{-Cat}^{T'}$$

of categories commuting with the forgetful functors into $\mathcal{G}V$.

PROOF. (1) $\Leftrightarrow$ (2): The basic idea of this proof is to adapt the discussion of [7] section 1 replacing one of the monads by a multitensor. Suppose maps τ_{X_i} are given which make T into a monoidal monad. Define $u'_X = \eta_{E_1 X} u_X$ and $\sigma'_{X_{ij}}$ to be given by the composite

$$T E_i T E_j X_{ij} \xrightarrow{T \tau E} T^2 E_i E_j X_{ij} \xrightarrow{\mu \sigma} T E_{ij} X_{ij}.$$

The axioms exhibiting (TE, u', σ') as a multitensor, $\eta_E : E \rightarrow TE$ as a multitensor map, $Tu : T \rightarrow TE_1$ as a monad map, and $\sigma'(Tu \eta E_i) = \text{id}$ follow easily from the multitensor axioms on E , the monad axioms on T and the monoidal monad coherence axioms. Conversely given the data σ' as in (2) one defines the monoidal monad coherence τ as the composite

$$E T_i \xrightarrow{\eta E_i T u} T E_i T E_1 \xrightarrow{\sigma'} T E_i.$$

The axioms involving τ , η and u are verified easily.

In order to verify the other two axioms one must first observe that

$$(7) \quad \begin{array}{ccc} TE_i E_j & \xrightarrow{T E_i \eta_j E_j} & TE_i TE_j \\ & \searrow T\sigma & \downarrow \sigma' \\ & & TE_{ij} \end{array} \quad \begin{array}{ccc} T^2 E_i & \xrightarrow{Tu TE_i} & TE_1 TE_i \\ & \searrow \mu_i^{E_i} & \downarrow \sigma' \\ & & TE_i \end{array}$$

commutes. One witnesses the commutativity of the triangle on the left from

$$\begin{array}{ccccccc} TE_i E_j & \xrightarrow{Tu \eta_i E_j E_j} & TE_1 TE_i E_j & \xrightarrow{\sigma'_j E_j} & TE_i E_j & \xrightarrow{T E_i \eta_j E_j} & TE_i TE_j \\ \swarrow T\sigma & \searrow Tu E_i E_j & \searrow TE_1 \eta_i E_j E_j & \searrow TE_1 TE_i \eta_j E_j & \searrow \sigma'_j TE_j & \searrow TE_i \eta_j E_j & \\ TE_{ij} & & TE_1 E_{ij} & \xrightarrow{TE_1 \eta_i E_j E_j} & TE_1 TE_i TE_j & \xrightarrow{\sigma'_j TE_j} & TE_i TE_j \\ \searrow Tu E_{ij} & \swarrow TE_1 \sigma & \swarrow TE_1 \sigma' & \swarrow \sigma' & \swarrow \sigma' & \swarrow \sigma' & \\ TE_1 E_{ij} & \xrightarrow{TE_1 \eta_{ij} E_j} & TE_1 TE_{ij} & \xrightarrow{\sigma'} & TE_{ij} & & \end{array}$$

using also $\sigma'(Tu \eta_i E_j) = \text{id}$, and one witnesses the commutativity of the triangle on the right of (7) from

$$\begin{array}{ccccccc} T^2 E_i & \xrightarrow{T^2 u \eta_i E_i} & T^2 E_1 TE_i & \xrightarrow{T\sigma'} & T^2 E_i & \xrightarrow{Tu TE_i} & TE_1 TE_i \\ \swarrow \mu_i^{E_i} & \searrow T^2 \eta_i E_i & \searrow Tu TE_1 TE_i & \searrow TE_1 \sigma' & \searrow \sigma' & \searrow \sigma' & \\ TE_i & & T^3 E_i & \xrightarrow{Tu Tu TE_i} & TE_1 TE_1 TE_i & \xrightarrow{TE_1 \sigma'} & TE_1 TE_i \\ \searrow T \eta_i E_i & \swarrow \mu T E_i E_1 & \swarrow \sigma' TE_i & \swarrow \sigma' & \swarrow \sigma' & \swarrow \sigma' & \\ T^2 E_i & \xrightarrow{Tu TE_i} & TE_1 TE_i & \xrightarrow{\sigma'} & TE_i & & \end{array}$$

and $\sigma'(Tu \eta_i E_j) = \text{id}$. Second, one observes that

$$(8) \quad \begin{array}{ccc} T^2 E_i TE_j & \xrightarrow{T\sigma'} & T^2 E_{ij} \\ \mu_i^{E_i} TE_j \downarrow & & \downarrow \mu_{ij}^{E_i} \\ TE_i TE_j & \xrightarrow{\sigma'} & TE_{ij} \end{array} \quad \begin{array}{ccc} TE_i TE_j E_k & \xrightarrow{\sigma'_k E_k} & TE_{ij} E_k \\ TE_i T\sigma \downarrow & & \downarrow T\sigma \\ TE_i TE_{jk} & \xrightarrow{\sigma'} & TE_{ijk} \end{array}$$

commute, but these identities are easily witnessed from

$$\begin{array}{ccc} T^2 E_i TE_j & \xrightarrow{T\sigma'} & T^2 E_{ij} \\ Tu TE_i TE_j \downarrow & & \downarrow Tu TE_{ij} \\ TE_1 TE_i TE_j & \xrightarrow{TE_1 \sigma'} & TE_1 TE_{ij} \\ \sigma'_j TE_j \downarrow & & \downarrow \sigma' \\ TE_i TE_j & \xrightarrow{\sigma'} & TE_{ij} \end{array} \quad \begin{array}{ccc} TE_i TE_j E_k & \xrightarrow{\sigma'_k E_k} & TE_{ij} E_k \\ TE_i TE_j E_k \downarrow & & \downarrow TE_{ij} E_k \\ TE_i TE_j TE_k & \xrightarrow{\sigma'_k TE_k} & TE_{ij} TE_k \\ TE_i \sigma' \downarrow & & \downarrow \sigma' \\ TE_i TE_{jk} & \xrightarrow{\sigma'} & TE_{ijk} \end{array}$$

and (7).

With (8) now verified we now proceed to the verification of the other axioms for τ . The axiom expressing the compatibility between τ and σ is verified in

$$\begin{array}{ccccccc}
 E E T_{ij} & \xrightarrow{E_i \eta_j E_j T u} & E T E_j T E_1 & \xrightarrow{E_i \sigma'} & E T E_j & \xrightarrow{\eta_i E_j T u E_j} & T E T E_1 E_j \\
 \downarrow \sigma T & \searrow \eta_i E_i \eta_j T u & \downarrow \eta_i E_i T E_j T E_1 & & \downarrow \eta_i E_i T E_j & \swarrow T E_i T \sigma & \downarrow \sigma' E_j \\
 & & T E T E_j T E_1 & \xrightarrow{T E_i \sigma'} & T E T E_j & & T E E_j \\
 & & \searrow \sigma' T E_1 & & \searrow \sigma' & & \downarrow T \sigma \\
 E T_{ij} & \xrightarrow{\eta_{ij} E T u} & T E T E_1 & \xrightarrow{\sigma'} & T E_{ij} & &
 \end{array}$$

and the axiom expressing the compatibility between τ and μ is verified in

$$\begin{array}{ccccccc}
 E T^2_i & \xrightarrow{\eta_i E_i T u T} & T E T E_1 T_i & \xrightarrow{\sigma' T} & T E T_i & \xrightarrow{T \eta_i E_i T u} & T^2 E T E_1_i \\
 \downarrow E_i \mu & \searrow \eta_i E_i T u T u & \downarrow T E_i T E_1 T u & & \downarrow T E_i T u & \swarrow \mu E_i T E_1 & \downarrow T \sigma' \\
 & & T E T E_1 T E_1 & \xrightarrow{\sigma' T E_1} & T E T E_1 & & T^2 E_i \\
 & & \searrow T E_i \sigma' & & \searrow \sigma' & & \downarrow \mu E_i \\
 E T_i & \xrightarrow{\eta E_1 T u} & T E T E_1 & \xrightarrow{\sigma'} & T E_i & &
 \end{array}$$

It follows immediately from the unit laws of T and E , that the composite function $(1) \rightarrow (2) \rightarrow (1)$ via the above constructions is the identity. It is the commutativity of the outside of

$$\begin{array}{ccccc}
 & & T^2 E T E_1 E_j & \xrightarrow{T \sigma' E_j} & T^2 E E_{ij} \\
 & \nearrow T \eta_i E_i T u E_j & \downarrow \mu E_i T E_1 E_j & & \downarrow \mu E_{ij} \\
 T E T E_{ij} & & T E T E_1 E_j & \xrightarrow{\sigma' E_j} & T E E_{ij} \\
 & \searrow 1 & \downarrow T E_i T \sigma & & \downarrow T \sigma \\
 & & T E T E_{ij} & \xrightarrow{\sigma'} & T E_{ij}
 \end{array}
 \quad \mu \sigma$$

that says that $(2) \rightarrow (1) \rightarrow (2)$ is the identity.

(1) $\Leftrightarrow$ (3): A monoidal monad on (V, E) is a monad on (V, E) in the 2-category **DISTMULT**. By [24] to give a monad S on $\mathcal{G}V$ over **Set** and a distributive law of

S over ΓE , is to give a monad on $(\mathcal{G}V, \Gamma E)$ in $\text{MND}(\mathbf{CAT}/\mathbf{Set})$. Moreover such distributive laws, for the case where $S = \mathcal{G}T$ for some monad T on V , are exactly monads in $\mathcal{G}\text{-MND}$. Thus applying Ψ to monads gives the desired bijection.

(3) $\Leftrightarrow$ (4): By the usual theory of monad distributive laws and since $\mathcal{G}V^{\Gamma E} \cong E\text{-Cat}$ over $\mathcal{G}V$ by proposition(26).

One can readily unpack the lifted monad T' in terms of the monoidal coherence data using the details of the proof of proposition(26) which explain how to regard an E -category as a ΓE -algebra. Let $X \in E\text{-Cat}$ and as in section(4.1) write $\kappa_{x_i} : \mathbb{E}_i X(x_{i-1}, x_i) \rightarrow X(x_0, x_n)$ for the composition maps. Then since T' is a lifting of $\mathcal{G}T$, $T'X$ must have underlying V -graph $\mathcal{G}T(X)$, which has the same objects as X and homs given by $(T'X)(a, b) = T(X(a, b))$. The composition map $\kappa'_{x_i} : \mathbb{E}_i T'X(x_{i-1}, x_i) \rightarrow T'X(x_0, x_n)$ is given by the composite

$$\mathbb{E}_i T'X(x_{i-1}, x_i) \xrightarrow{\tau} T \mathbb{E}_i X(x_{i-1}, x_i) \xrightarrow{T\kappa_{x_i}} TX(x_0, x_n)$$

Consider $Z \in \mathcal{G}V$. To endow Z with the structure of a TE -category is to give maps

$$\kappa_{z_i} : T \mathbb{E}_i Z(z_{i-1}, z_i) \rightarrow Z(z_0, z_n)$$

satisfying the usual axioms. But by precomposing these with the unit for T gives the compositions for an E -category structure on Z , and by the above explicit description of T' , one may readily verify that the remaining structure is exactly that of a T' -algebra structure. Similarly given V -graph map $f : Z \rightarrow Z'$ between TE -categories, one may readily verify that f is a TE -functor iff it is an E -functor and a T' -algebra map. Thus we have the object and arrow maps of the required isomorphism $(TE)\text{-Cat} \cong E\text{-Cat}^{T'}$. $\square$

All aspects of the above result apply to the familiar examples of monoidal monads on $\mathbf{Set}$ regarded as monoidal via its cartesian product, since $\times$ preserves all colimits in each variable and so is certainly distributive. These familiar examples include: the pointed set monad, the covariant power set monad, the monad obtained from a commutative ring R by taking R -linear combinations. In [27] a tensor product on $\mathcal{G}V$ is provided, under very slight conditions on V , with respect to which any monad on $\mathcal{G}V$ over $\mathbf{Set}$ is (symmetric) monoidal, giving many examples relevant to higher category theory.

6.2. Distributing a multitensor over a monad. In a completely analogous fashion one may also regard opmonoidal monads as distributive laws. Once again let (V, E, u, σ) be a lax monoidal category. An *opmonoidal monad* on (V, E) is a monad on (V, E) in the 2-category of lax monoidal categories, oplax monoidal functors and monoidal natural transformations, which amounts to a monad (T, η, μ) on V together with coherence maps

$$\tau_{X_i} : T \mathbb{E}_i X_i \rightarrow \mathbb{E}_i T X_i$$

such that

$$\begin{array}{ccc}
 T & \xrightarrow{Tu} & TE_1 \\
 uT \searrow & & \swarrow \tau \\
 & E_1T &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 TEE_{ij} & \xrightarrow{\tau E_j} & ET E_j & \xrightarrow{E_i \tau} & EE_iT \\
 T\sigma \searrow & & & & \swarrow \sigma T \\
 & TE_{ij} & \xrightarrow{\tau} & ET_{ij} &
 \end{array}$$

and

$$\begin{array}{ccc}
 E_1 & \xrightarrow{\eta E_1} & TE_1 \\
 E_1\eta \searrow & & \swarrow \tau \\
 & E_1T &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 T^2E_i & \xrightarrow{T\tau} & TE_iT & \xrightarrow{\tau T} & E_iT^2 \\
 \mu E_i \searrow & & & & \swarrow E_i\mu \\
 & TE_i & \xrightarrow{\tau} & ET_i &
 \end{array}$$

commute. Recalling that M is our notation for the monoid monad on $\mathbf{CAT}$, we shall use the alternative notations

$$EM(T)(X_1, \dots, X_n) = \underset{i}{E}TX_i$$

interchangeably as convenience dictates.

Given a multitensor (E, u, σ) on a category V and a monad (T, η, μ) on V , a *lifting of E to V^T* is defined to be a multitensor (E', u', σ') on V^T such that $U^TE' = EM(U^T)$, $U^Tu' = uU^T$ and $U^T\sigma' = \sigma M^2(U^T)$, where we recall that $U^T : V^T \rightarrow V$ is the forgetful functor. In more explicit terms to give such a lifting is to give maps

$$a_{x_i} : T \underset{i}{E} X_i \rightarrow \underset{i}{E} X_i$$

in V for all sequences $((X_1, x_1), \dots, (X_n, x_n))$ of T -algebras, such that these maps satisfy the axioms making $(\underset{i}{E} X_i, a_{x_i})$ T -algebras, and with respect to these structures, $\underset{i}{E} f_i$ is a morphism of T -algebras for any sequence $f_i : (X_i, x_i) \rightarrow (Y_i, y_i)$ of T -algebra maps, and moreover, u and σ are T -algebra morphisms.

Theorem 49. *Let V be a category, (E, u, σ) be a multitensor on V and (T, η, μ) be a monad on V . Then there is a bijection between the following types of data:*

- (1) *Morphisms $\tau_{X_i} : T \underset{i}{E} X_i \rightarrow \underset{i}{E} TX_i$ of V providing the coherences making T into a opmonoidal monad.*
- (2) *Morphisms $\sigma'_{X_{ij}} : \underset{i}{E} T \underset{j}{E} TX_{ij} \rightarrow \underset{ij}{E} TX_{ij}$ of V providing the substitutions for a multitensor $(EM(T), u', \sigma')$ where $u'_X = u_{TX} \eta_X$, $E\eta : E \rightarrow EM(T)$ is a multitensor map, $uT : T \rightarrow E_1T$ is a monad map, and the composite*

$$\underset{i}{E} T \xrightarrow{\underset{i}{E} \eta u T} \underset{i}{E} T E_1 T \xrightarrow{\sigma'} \underset{i}{E} T$$

is the identity.

- (3) *Liftings of the multitensor E to a multitensor E' on V^T .*

and for any given instance of such data one has an isomorphism

$$EM(T)\text{-Cat} \cong E'\text{-Cat}$$

of categories commuting with the forgetful functors into $\mathcal{GV}$. If in addition V has coproducts, T preserves them and E is distributive, then the data (1)-(2) are also in bijection with

- (4) Identity on object morphisms $\lambda_X : \mathcal{G}(T)\Gamma(E)(X) \rightarrow \Gamma(E)\mathcal{G}(T)(X)$ of $\mathcal{GV}$ providing a monad distributive law of $\Gamma(E)$ over $\mathcal{G}(T)$.

PROOF. (1) $\Leftrightarrow$ (2): This is completely analogous to the bijection between (1) and (2) in theorem(48).

(1) $\Leftrightarrow$ (3): Suppose that opmonoidal monad coherence data τ_{X_i} is given. Then for a given sequence of T -algebras $((X_1, x_1), \dots, (X_n, x_n))$ we define a_{x_i} to be given by the composite

$$T E X_i \xrightarrow{\tau} E T X_i \xrightarrow{E x_i} E X_i.$$

The verifications that these maps satisfy the T -algebra axioms, and that with respect to these structures u and σ are T -algebra morphisms are straight forward. Conversely given a lifting of E to V^T we construct, for a given sequence $(X_1, \dots, X_n)$ of objects of V , the coherence map τ_{X_i} as the composite

$$T E X_i \xrightarrow{T E \eta} T E T X_i \xrightarrow{a_{\mu X_i}} E X_i.$$

The axioms expressing the compatibilities between τ with u , η and μ are all routinely verified. The axiom for the compatibility of τ with σ is witnessed in

$$\begin{array}{ccccc}
 T E E X_{ij} & \xrightarrow{T E \eta E_j} & T E T E X_{ij} & \xrightarrow{a_{\mu E_j X_{ij}}} & E T E X_{ij} \\
 \downarrow T \sigma & \searrow T E E \eta & \downarrow T E T E \eta & & \downarrow E T E \eta \\
 T E X_{ij} & & T E T E T X_{ij} & \xrightarrow{a_{\mu E_j T X_{ij}}} & E T E T X_{ij} \\
 \downarrow T E \eta & \nearrow T E \eta E_j T & \downarrow T E a_{\mu X_{ij}} & & \downarrow E a_{\mu X_{ij}} \\
 T E T X_{ij} & \xrightarrow{1} & T E E T X_{ij} & \xrightarrow{a_{a \mu X_{ij}}} & E E T X_{ij} \\
 \downarrow a_{\mu X_{ij}} & & \downarrow \sigma T & & \\
 E T X_{ij} & & E T X_{ij} & &
 \end{array}$$

(I) (II) (III)

in which the unlabelled regions commute for obvious reasons. Region (III) commutes since σ is a T -algebra map. Since the $a_{\mu X_{ij}}$ are T -algebra maps so is $E a_{\mu X_{ij}}$ and so region (II) commutes. The morphisms $T E \eta_j$ are T -algebra maps and so $E T E \eta$ are T -algebra maps whence region (I) commutes also. Thus we have established functions that turn opmonoidal coherence data into liftings and vice versa, and the verification that these are inverse to each other is straight forward.

The isomorphism $E'\text{-Cat} \cong (EM(T))\text{-Cat}$ over $\mathcal{GV}$: To give $X \in \mathcal{GV}$ the structure of an $EM(T)$ -category is to give morphisms

$$a_{x_i} : \mathbb{E}_i T(X(x_{i-1}, x_i)) \rightarrow X(x_0, x_n)$$

in V for each sequence $(x_0, \dots, x_n)$ of objects of X , such that all diagrams of the form

$$\begin{array}{ccc} X(x_0, x_1) & \xrightarrow{\eta} & TX(x_0, x_1) \\ \downarrow 1 & & \downarrow uT \\ X(x_0, x_1) & \xleftarrow{a_{x_0, x_1}} & E_1 TX(x_0, x_1) \end{array} \quad \begin{array}{ccc} \mathbb{E}_i T \mathbb{E}_j TX(x_{ij-1}, x_{ij}) & \xrightarrow{\mathbb{E}_i \tau T} & \mathbb{E}_i \mathbb{E}_j T^2 X(x_{ij-1}, x_{ij}) \\ \downarrow \mathbb{E}_i T a_{x_{ij}} & & \downarrow \sigma \mu \\ \mathbb{E}_i TX(x_{i-1}, x_i) & & \mathbb{E}_{ij} TX(x_{ij-1}, x_{ij}) \\ & \searrow a_{a_i} & \swarrow a_{x_{ij}} \\ & X(x_0, x_n) & \end{array}$$

commute. On the hand to give X the structure of an E' -category is to give maps


$$b_{x_0, x_1} : TX(x_0, x_1) \rightarrow X(x_0, x_1) \quad c_{x_i} : \mathbb{E}_i X(x_{i-1}, x_i) \rightarrow X(x_0, x_n)$$

such that the b_{x_0, x_1} satisfy the T -algebra axioms, the c_{x_i} satisfy the E -category axioms, and moreover the c_{x_i} are T -algebra morphisms with respect to the T -algebra structures given by the b_{x_0, x_1} . Supposing that maps a_{x_i} are given as above one defines maps b_{x_0, x_1} and c_{x_i} as composites

$$\begin{array}{ccccc} TX(x_0, x_1) & \xrightarrow{uT} & E_1 TX(x_0, x_1) & \xrightarrow{a_{x_0, x_1}} & X(x_0, x_1) \\ \mathbb{E}_i X(x_{i-1}, x_i) & \xrightarrow{\mathbb{E}_i \eta} & \mathbb{E}_i TX(x_{i-1}, x_i) & \xrightarrow{a_{x_i}} & X(x_0, x_n). \end{array}$$

The T -algebra axioms for b_{x_0, x_1} and the E -category axioms for c_{x_i} are easily verified. That c_{x_i} is a T -algebra morphism is expressed by the commutativity of the outside

of the diagram



whose internal regions all clearly commute. Conversely given structure maps b_{x_0, x_1} and c_{x_i} as above, one constructs the a_{x_i} as the composites

$$E_i T X(x_{i-1}, x_i) \xrightarrow{E_i b_{x_{i-1}, x_i}} E_i X(x_{i-1}, x_i) \xrightarrow{c_{x_i}} X(x_0, x_n).$$

The verification that these maps satisfy the axioms on the a_{x_i} described above is straight forward. It is also straight forward to check that these constructions give a bijection between $EM(T)$ -category structures and E' -category structures on X , and moreover that this can be extended to maps giving the required isomorphism of categories over $\mathcal{GV}$.

(1) $\Leftrightarrow$ (4): This bijection is given in basically the same way as that for (1) $\Leftrightarrow$ (3) in theorem(48), using the 2-functor Ψ' instead of Ψ . $\square$

Remark 50. Note in particular that, in the context of theorem(49), when V has coproducts, E is distributive and T preserves coproducts, then the composite multitensor $EM(T)$ is clearly distributive. Moreover since $U^T E' = EM(U^T)$ and U^T

creates coproducts, the lifted multitensor E' is also distributive. Thus by theorem(49) and proposition(26) one in fact has isomorphisms

$$\mathcal{G}(V)^{\Gamma(EM(T))} \cong EM(T)\text{-Cat} \cong \mathcal{G}(V)^{\Gamma(E)\mathcal{G}(T)} \cong E'\text{-Cat} \cong \mathcal{G}(V^T)^{\Gamma E'}$$

over $\mathcal{G}V$. Either by a direct verification, or by applying structure-semantics² to $\mathcal{G}(V)^{\Gamma(EM(T))} \cong \mathcal{G}(V)^{\Gamma(E)\mathcal{G}(T)}$, one has also an isomorphism

$$\Gamma(EM(T)) \cong \Gamma(E)\mathcal{G}(T)$$

of monads. Moreover the monad T may also be regarded as a multitensor, whose unary part is T and whose n -ary parts for $n \neq 1$ are constant at $\emptyset$, and then $\Gamma T \cong \mathcal{G}T$ as monads. Thus if E and T are λ -accessible, then so is $EM(T)$ by theorem(29)(2).

Remark 51. If moreover E and T are l.r.a and the coherences τ_{X_i} are cartesian natural in the X_i , then by the explicit description of the composite multitensor $EM(T)$ and theorem(29)(4), $EM(T)$ is also l.r.a.

Example 52. When V is a category with finite products and E is given by them, any monad T on V is canonically opmonoidal, with the coherences provided by the product preservation obstruction maps. The composite monad, whose tensor product is

$$EM(T)(X_1, \dots, X_n) = \prod_{1 \leq i \leq n} TX_i$$

was called $T^\times$ in [6]. Proposition(2.8) of [6], which says that

$$T^\times\text{-Cat} \cong (V^T, \times)\text{-Cat}$$

over $\mathcal{G}V$, follows by applying theorem(49) to this example. When T is l.r.a and V is distributive, the product obstruction maps for T are cartesian natural by lemma(2.15) of [26] and so by remark(51) $T^\times$ is l.r.a.

6.3. Strict n -category monads. One can consider the following inductively-defined sequence of monads

- Put $\mathcal{T}_{\leq 0}$ equal to the identity monad on **Set**.
- Given a monad $\mathcal{T}_{\leq n}$ on $\mathcal{G}^n\mathbf{Set}$, define the monad $\mathcal{T}_{\leq n+1} = \Gamma\mathcal{T}_{\leq n}^\times$ on $\mathcal{G}^{n+1}\mathbf{Set}$.

recalling that $\mathcal{G}^n\mathbf{Set}$ is the category of n -globular sets. By example(52) and proposition(26) it follows that $\mathcal{G}^n(\mathbf{Set})^{\mathcal{T}_{\leq n}}$ is the category of strict n -categories and strict n -functors between them. By remarks(50) and (51), and example(52), we recover the fundamental properties of these monads, that is that they are coproduct preserving, finitary and l.r.a. Moreover from this inductive description of $\mathcal{T}_{\leq n}$ and theorem(49) one recovers the distributive law

$$\mathcal{G}(\mathcal{T}_{\leq n})\Gamma(\Pi) \rightarrow \Gamma(\Pi)\mathcal{G}(\mathcal{T}_{\leq n})$$

for all n , between monads on $\mathcal{G}^n\mathbf{Set}$, with composite monad $\Gamma(\Pi)\mathcal{G}(\mathcal{T}_{\leq n}) = \mathcal{T}_{\leq (n+1)}$, as witnessed in [10].

²This is the well-known fact due to Lawvere that for any category $\mathcal{E}$, the canonical functor

$$\mathbf{Mnd}(\mathcal{E})^{\text{op}} \rightarrow \mathbf{CAT}/\mathcal{E} \quad T \mapsto U : \mathcal{E}^T \rightarrow \mathcal{E}$$

with object map indicated is fully faithful (see [24] for a proof). In particular this implies that for monads S and T on $\mathcal{E}$, an isomorphism $\mathcal{E}^T \cong \mathcal{E}^S$ over $\mathcal{E}$ is the same thing as a monad isomorphism $S \cong T$.

One could just as well start with any locally finitely c-presentable V in place of **Set**, giving monads whose algebras are V -enriched strict n -categories. This enrichment gives objects in V of n -cells between parallel pairs of $(n-1)$ -cells. By the same arguments these monads are also coproduct preserving, finitary and l.r.a., and moreover give rise to distributive laws in the same way.

7. Contractible operads and multitensors

7.1. The basic correspondence between operads and multitensors.

Recall [6] that just as one can define T -operads for a cartesian monad T , one also has a notion of E -multitensor for any cartesian multitensor E . For (V, E) a cartesian multitensor, one defines an E -multitensor to consist of another multitensor A on V together with a natural transformation $\alpha : A \rightarrow E$ which is cartesian natural and compatible with the multitensor structures.

For V a category with pullbacks and T a cartesian monad on $\mathcal{G}V$ over **Set**, a T -operad $\alpha : A \rightarrow T$ over **Set** may be regarded as either a monad functor $(1_{\mathcal{G}V}, \alpha) : (\mathcal{G}V, T) \rightarrow (\mathcal{G}V, A)$ over **Set** whose 1-cell datum is an identity and 2-cell datum is a cartesian transformation, or equally well as a monad opfunctor $(1_{\mathcal{G}V}, \alpha) : (\mathcal{G}V, A) \rightarrow (\mathcal{G}V, T)$ whose 1-cell datum is an identity and 2-cell datum is cartesian. Similarly, an E -multitensor may be regarded as either a lax monoidal functor $(1_V, \alpha) : (V, E) \rightarrow (V, A)$ over **Set** whose 1-cell datum is an identity and 2-cell datum is a cartesian transformation, or equally well as an oplax monoidal functor $(1_V, \alpha) : (V, A) \rightarrow (V, E)$ whose 1-cell datum is an identity and 2-cell datum is cartesian.

We denote by $T\text{-Op}_0$ the category of T -operads over **Set** and their morphisms. A morphism from $\alpha : A \rightarrow T$ to $\beta : B \rightarrow T$ is just a monad morphism $\gamma : A \rightarrow B$ such that $\alpha = \beta\gamma$. It follows that γ is itself cartesian and over **Set**. Thus a T -operad morphism may be regarded either as a monad functor $(1_V, \gamma) : (V, T) \rightarrow (V, A)$ or a monad opfunctor $(1_V, \gamma) : (V, A) \rightarrow (V, T)$. Similarly one has the category $E\text{-Mult}$ of E -multitensors and their morphisms, with a morphism from $\alpha : A \rightarrow E$ to $\beta : B \rightarrow E$ being multitensor map over E . As with operad morphisms, morphisms of E -multitensors are reexpressable either as lax monoidal functors under (V, E) or as oplax monoidal functors over (V, E) . Thus by applying either Γ or Γ' to E -multitensors and their morphisms, one obtains a functor

$$\Gamma_E : E\text{-Mult} \rightarrow \Gamma E\text{-Op}_0.$$

By lemma(47), theorem(42) and proposition(46), Γ_E is essentially surjective on objects. By proposition(44) it is fully faithful, and so we have obtained

Theorem 53. *Let V be left extensive. Then Γ_E gives an equivalence of categories*

$$E\text{-Mult} \simeq \Gamma E\text{-Op}_0.$$

In the case where $E = \mathcal{T}_{\leq n}^\times$ we recover the first equivalence of corollary(7.10) and of corollary(8.3) of [6].

7.2. Trivial Fibrations. Let V be a category and $\mathcal{I}$ a class of maps in V . Denote by $\mathcal{I}^\uparrow$ the class of maps in V that have the right lifting property with respect to all the maps in $\mathcal{I}$. That is to say, $f : X \rightarrow Y$ is in $\mathcal{I}^\uparrow$ iff for every $i : S \rightarrow B$ in $\mathcal{I}$,

α and β such that the outside of

$$\begin{array}{ccc} S & \xrightarrow{\alpha} & X \\ i \downarrow & \nearrow \gamma & \downarrow f \\ B & \xrightarrow{\beta} & Y \end{array}$$

commutes, then there is a γ as indicated such that $f\gamma=\beta$ and $\gamma i = \alpha$. An $f \in \mathcal{I}^\uparrow$ is called a *trivial $\mathcal{I}$ -fibration*. The basic facts about $\mathcal{I}^\uparrow$ that we shall use are summarised in

Lemma 54. *Let V be a category, $\mathcal{I}$ a class of maps in V , J a set and*

$$(f_j : X_j \rightarrow Y_j \mid j \in J)$$

a family of maps in V .

- (1) $\mathcal{I}^\uparrow$ is closed under composition and retracts.
- (2) If V has products and each of the f_j is a trivial $\mathcal{I}$ -fibration, then

$$\prod_j f_j : \prod_j X_j \rightarrow \prod_j Y_j$$

is also a trivial $\mathcal{I}$ -fibration.

- (3) The pullback of a trivial $\mathcal{I}$ -fibration along any map is a trivial $\mathcal{I}$ -fibration.
- (4) If V is extensive and $\coprod_j f_j$ is a trivial $\mathcal{I}$ -fibration, then each of the f_j is a trivial fibration.
- (5) If V is extensive, the codomains of maps in $\mathcal{I}$ are connected and each of the f_j is a trivial $\mathcal{I}$ -fibration, then $\coprod_j f_j$ is a trivial $\mathcal{I}$ -fibration.

PROOF. (1)-(3) is standard. If V is extensive then the squares

$$\begin{array}{ccc} X_j & \longrightarrow & \coprod_j X_j \\ f_j \downarrow & & \downarrow \coprod_j f_j \\ Y_j & \longrightarrow & \coprod_j Y_j \end{array}$$

whose horizontal arrows are the coproduct injections are pullbacks, and so (4) follows by the pullback stability of trivial $\mathcal{I}$ -fibrations. As for (5) note that for $i : S \rightarrow B$ in $\mathcal{I}$, the connectedness of B ensures that any square as indicated on the left

$$\begin{array}{ccc} S & \longrightarrow & \coprod_j X_j \\ i \downarrow & & \downarrow \coprod_j f_j \\ B & \longrightarrow & \coprod_j Y_j \end{array} \qquad \begin{array}{ccc} S & \longrightarrow & X_j \\ i \downarrow & & \downarrow f_j \\ B & \longrightarrow & Y_j \end{array}$$

factors through a unique component as indicated on the right, enabling one to induce the desired filler. $\square$

Definition 55. Let $F, G : W \rightarrow V$ be functors and $\mathcal{I}$ be a class of maps in V . A natural transformation $\phi : F \Rightarrow G$ is a *trivial $\mathcal{I}$ -fibration* when its components are trivial $\mathcal{I}$ -fibrations.

Note that since trivial $\mathcal{I}$ -fibrations in V are pullback stable, this reduces, in the case where W has a terminal object 1 and ϕ is cartesian, to the map $\phi_1 : F1 \rightarrow G1$ being a trivial $\mathcal{I}$ -fibration.

Given a category V with an initial object, and a class of maps $\mathcal{I}$ in V , we denote by $\mathcal{I}^+$ the class of maps in $\mathcal{G}V$ containing the maps³

$$\emptyset \rightarrow 0 \quad (i) : (S) \rightarrow (B)$$

where $i \in \mathcal{I}$. The proof of the following lemma is trivial.

Lemma 56. *Let V be a category with an initial object and $\mathcal{I}$ a class of maps in V . Then $f : X \rightarrow Y$ is a trivial $\mathcal{I}^+$ -fibration iff it is surjective on objects and all its hom maps are trivial $\mathcal{I}$ -fibrations.*

In particular starting with $V = \widehat{\mathbb{G}}$ the category of globular sets and $\mathcal{I}_{-1}$ the empty class of maps, one generates a sequence of classes of maps $\mathcal{I}_n$ of globular sets by induction on n by the formula $\mathcal{I}_{n+1} = (\mathcal{I}_n)^+$ since $\mathcal{G}(\widehat{\mathbb{G}})$ may be identified with $\widehat{\mathbb{G}}$, and moreover one has inclusions $\mathcal{I}_n \subset \mathcal{I}_{n+1}$. More explicitly, the set $\mathcal{I}_n$ consists of $(n+1)$ maps: for $0 \leq k \leq n$ one has the inclusion $\partial k \hookrightarrow k$, where k here denotes the representable globular set, that is the “ k -globe”, and ∂k is the k -globe with its unique k -cell removed. One defines $\mathcal{I}_{\leq \infty}$ to be the union of the $\mathcal{I}_n$ ’s. Note that by definition $\mathcal{I}_{\leq \infty} = \mathcal{I}_{\leq \infty}^+$.

There is another version of the induction just described to produce, for each $n \in \mathbb{N}$, a class $\mathcal{I}_{\leq n}$ of maps of $\mathcal{G}^n(\mathbf{Set})$. The set $\mathcal{I}_{\leq 0}$ consists of the functions

$$\emptyset \rightarrow 0 \quad 0 + 0 \rightarrow 0,$$

so $\mathcal{I}_{\leq 0}^+$ is the class of bijective functions. For $n \in \mathbb{N}$, $\mathcal{I}_{\leq n+1} = \mathcal{I}_{\leq n}^+$. As maps of globular sets, the class $\mathcal{I}_{\leq n}$ consists of all the maps of $\mathcal{I}_n$ together with the unique map $\partial(n+1) \rightarrow n$.

Definition 57. Let $0 \leq n \leq \infty$. An n -operad⁴ $\alpha : A \rightarrow \mathcal{T}_{\leq n}$ is *contractible* when it is a trivial $\mathcal{I}_{\leq n}$ -fibration. An n -multitensor $\varepsilon : E \rightarrow \mathcal{T}_{\leq n}^\times$ is *contractible* when it is a trivial $\mathcal{I}_{\leq n}$ -fibration.

By the preceding two lemmas, an $(n+1)$ -operad $\alpha : A \rightarrow \mathcal{T}_{\leq n+1}$ over $\mathbf{Set}$ is contractible iff the hom maps of α_1 are trivial $\mathcal{I}_{\leq n}$ -fibrations.

7.3. Contractible operads versus contractible multitensors. As one would expect an $\mathcal{T}_{\leq n+1}$ -operad over $\mathbf{Set}$ is contractible iff its associated $\mathcal{T}_{\leq n}^\times$ -multitensor is contractible. This fact has quite a general explanation.

Proposition 58. *Let $(H, \psi) : (V, E) \rightarrow (W, F)$ be a lax monoidal functor between distributive lax monoidal categories, and $\mathcal{I}$ a class of maps in W . Suppose that W is extensive, H preserves coproducts and the codomains of maps in $\mathcal{I}$ are connected. Then the following statements are equivalent*

- (1) ψ is a trivial $\mathcal{I}$ -fibration.
- (2) $\Gamma\psi$ is a trivial $\mathcal{I}^+$ -fibration.

³Recall that 0 is the V -graph with one object whose only hom is initial, or in other words the representing object of the functor $\mathcal{G}V \rightarrow \mathbf{Set}$ which sends a V -graph to its set of objects.

⁴The monad $\mathcal{T}_{\leq \infty}$ on globular sets is usually just denoted as $\mathcal{T}$: it is the monad whose algebras are strict ω -categories.

PROOF. For each $X \in \mathcal{GV}$ the component $\{\Gamma\psi\}_X$ is the identity on objects and for $a, b \in X_0$, the corresponding hom map is obtained as the composite of

$$\coprod_{a=x_0, \dots, x_n=b} \psi : \coprod_{x_0, \dots, x_n} F_i H X(x_{i-1} x_i) \rightarrow \coprod_{x_0, \dots, x_n} H E_i X(x_{i-1} x_i)$$

and the canonical isomorphism that witnesses the fact that H preserves coproducts. In particular note that for any sequence $(Z_1, \dots, Z_n)$ of objects of V , regarded as V -graph in the usual way, one has

$$\{\Gamma\psi\}_{(Z_1, \dots, Z_n)} = \psi_{Z_1, \dots, Z_n}.$$

Thus (1) $\Leftrightarrow$ (2) follows from lemmas(54) and (56). $\square$

Corollary 59. *Let $0 \leq n \leq \infty$, $\alpha : A \rightarrow \mathcal{T}_{\leq n+1}$ be a $\mathcal{T}_{\leq n+1}$ -operad over **Set** and $\varepsilon : E \rightarrow \mathcal{T}_{\leq n}^\times$ be the corresponding $\mathcal{T}_{\leq n}^\times$ -multitensor. TFSAE:*

- (1) $\alpha : A \rightarrow \mathcal{T}_{\leq n+1}$ is contractible.
- (2) $\varepsilon : E \rightarrow \mathcal{T}_{\leq n}^\times$ is contractible.

PROOF. By induction one may easily establish that the codomains of the maps in any of the classes: $\mathcal{I}_n, \mathcal{I}_{\leq n}, \mathcal{I}_{\leq \infty}$ are connected so that proposition(58) may be applied. $\square$

8. Trimble's construction

In this section we exhibit Cheng's analysis [11] of Trimble's definition as fitting within our framework.

8.1. Topological preliminaries. Given a topological space X and points a and b therein, one may define the topological space $X(a, b)$ of paths in X from a to b at a high degree of generality. In recalling the details let us denote by **Top** a category of "spaces" which is complete, cocomplete and cartesian closed. We shall write 1 for the terminal object. We shall furthermore assume that **Top** comes equipped with a bipointed object I playing the role of the interval. A conventional choice for **Top** is the category of compactly generated Hausdorff spaces with its usual interval, although there are many other alternatives which would do just as well from the point of view of homotopy theory.

Let us denote by σX the suspension of X , which can be defined as the pushout

$$\begin{array}{ccc} X+X & \longrightarrow & I \times X \\ \downarrow & & \downarrow \\ 1+1 & \xrightarrow{\quad \sqcup \quad} & \sigma X. \end{array}$$

Writing **Top_•** for the category of bipointed spaces, that is to say the coslice $1+1/\mathbf{Top}$, the above definition exhibits the suspension construction as a functor

$$\sigma : \mathbf{Top} \rightarrow \mathbf{Top}_\bullet.$$

Applying σ successively to the inclusion of the empty space into the point, one obtains the inclusions of the $(n-1)$ -sphere into the n -disk for all $n \in \mathbb{N}$, and its right adjoint

$$h : \mathbf{Top}_\bullet \rightarrow \mathbf{Top}$$

is the functor which sends the bipointed space (a, X, b) , to the space $X(a, b)$ of paths in X from a to b . This adjunction $\sigma \dashv h$ is easy to verify directly using the above elementary definition of $\sigma(X)$ as a pushout, and the pullback square

$$\begin{array}{ccc} X(a, b) & \xrightarrow{\quad} & X^I \\ \downarrow & \lrcorner & \downarrow X^i \\ 1 & \xrightarrow{(a, b)} & X^{1+1} \end{array}$$

where i is the inclusion of the boundary of I .

Thus to each space X one can associate a canonical topologically enriched graph whose homs are the path spaces of X . In section(3) we described explicitly the adjunction $(-)^{\times 2}_{\bullet} \dashv \mathcal{G}_1$ and in particular its unit η , from which one may readily verify that the assignment $X \mapsto PX$ is the object map of the composite

$$\mathbf{Top} \xrightarrow{\eta} \mathcal{G}(\mathbf{Top}_{\bullet}) \xrightarrow{\mathcal{G}h} \mathcal{G}\mathbf{Top}$$

and by proposition(14), the component η_f of the unit of this adjunction at $f : A \rightarrow \mathbf{Set}$ has a left adjoint when A is cocomplete. Since $\mathbf{Top}$ is cocomplete, h has left adjoint σ , and $\mathcal{G}$ is a 2-functor, P is a right adjoint.

Recall from section(4.1) that non-symmetric operads within braided monoidal categories may be regarded as multitensors, and that these are distributive when the tensor product is distributive. To say that a non-symmetric topological operad A acts on P is to say that P factors as

$$\mathbf{Top} \xrightarrow{P_A} A\text{-Cat} \xrightarrow{U^A} \mathcal{G}(\mathbf{Top})$$

The main example to keep in mind is the version of the little intervals operad recalled in [11] definition(1.1). As this A is a contractible non-symmetric operad, A -categories may be regarded as a model of A -infinity spaces. Since P is a right adjoint, P_A is also a right adjoint by the Dubuc adjoint triangle theorem.

8.2. Inductive construction. Let A be a non-symmetric topological operad which acts on P . Applying a product preserving functor

$$Q : \mathbf{Top} \rightarrow V$$

into a distributive category to the operad A in $\mathbf{Top}$, produces an operad QA in V . Moreover Q may be regarded as the underlying functor of a strong monoidal functor $(\mathbf{Top}, A) \rightarrow (V, QA)$ between lax monoidal categories. Applying Γ to this gives us a monad functor

$$(\mathcal{G}(\mathbf{Top}), \Gamma(A)) \rightarrow (\mathcal{G}V, \Gamma(QA))$$

with underlying functor $\mathcal{G}Q$, which amounts to giving a lifting $\overline{Q}$ as indicated in the commutative diagram

$$\begin{array}{ccccc} \mathbf{Top} & \xrightarrow{P_A} & A\text{-Cat} & \xrightarrow{\overline{Q}} & QA\text{-Cat} \\ & \searrow P & \downarrow & & \downarrow U^{QA} \\ & & \mathcal{G}(\mathbf{Top}) & \xrightarrow{\mathcal{G}(Q)} & \mathcal{G}V \end{array}$$

and so we have produced another product preserving functor

$$Q^{(+)} : \mathbf{Top} \rightarrow V^{(+)}$$

where $Q^{(+)} = \overline{Q}P_A$ and $V^{(+)} = QA\text{-Cat}$. The functor $\overline{Q}$ is product preserving since $\mathcal{G}(Q)$ is and U^{QA} creates products. The assignment

$$(Q, V) \mapsto (Q^{(+)}, V^{(+)})$$

in the case where A is as described in [11] definition(1.1), is the inductive process lying at the heart of the Trimble definition. In this definition one begins with the path components functor $\pi_0 : \mathbf{Top} \rightarrow \mathbf{Set}$ and defines the category $\mathbf{Trm}_0$ of “Trimble 0-categories” to be $\mathbf{Set}$. The induction is given by

$$(\mathbf{Trm}_{n+1}, \pi_{n+1}) := (\mathbf{Trm}_n^{(+)}, \pi_n^{(+)})$$

and so this definition constructs not only a notion of weak n -category but the product preserving π_n ’s to be regarded as assigning the fundamental n -groupoid to a space.

8.3. Operads for Trimble n -categories. In the context of a product preserving functor $Q : \mathbf{Top} \rightarrow V$ as above, suppose that W is a lextensive category, T is a coproduct preserving cartesian monad on W , and $\phi : S \rightarrow T$ is a T -operad. Suppose moreover that $V = W^S$. Then the operad/multitensor QA is a lifting of the operad/multitensor U^SQA , and so $QA\text{-Cat}$ may be identified with categories enriched in the multitensor on W whose tensor product is given by $(U^SQA)_n \times SX_1 \times \dots \times SX_n$ by theorem(49). But the composites

$$(9) \quad U^SQA(A)_n \times \prod_i SX_i \xrightarrow{\text{proj}} \prod_i SX_i \xrightarrow{\prod_i \phi_{X_i}} \prod_i TX_i$$

are the components of a cartesian multitensor map into $T^\times$. Thus by theorem(53) $(QA)\text{-Cat}$ is the category of algebras of a $\Gamma(T^\times)$ -operad over $\mathbf{Set}$. Thus by the inductive definition of $\mathbf{Trm}_n$ and of the monads $\mathcal{T}_{\leq n}$, $\mathbf{Trm}_n$ is the category of algebras of a $\mathcal{T}_{\leq n}$ -operad.

8.4. Contractibility of the Trimble operads. Let us denote by $\mathcal{J}$ the set of inclusions $S^{n-1} \rightarrow D^n$ of the n -sphere into the n -disk for $n \in \mathbb{N}$. Recall that these may all be obtained by successively applying the suspension functor σ to the inclusion of the empty space into the point. By definition the given topological operad A is contractible when for each n the unique map $A_n \rightarrow 1$ is in $\mathcal{J}^\uparrow$, and this is equivalent to saying that the cartesian multitensor map $A \rightarrow \prod$ is a trivial $\mathcal{J}$ -fibration. We shall write $U_n : \mathbf{Trm}_n \rightarrow \mathcal{G}^n \mathbf{Set}$ for the forgetful functor for each n .

Lemma 60. *If $f : X \rightarrow Y$ is a trivial $\mathcal{J}$ -fibration then*

- (1) $f_{a,b} : X(a,b) \rightarrow Y(fa,fb)$ is a trivial $\mathcal{J}$ -fibration for all $a, b \in X$.
- (2) $U_n \pi_n f$ is a trivial $\mathcal{I}_{\leq n}$ -fibration.

PROOF. (1): To give a commutative square as on the left in

$$\begin{array}{ccc} S^{n-1} & \longrightarrow & X(a,b) \\ \downarrow & & \downarrow \\ D^n & \longrightarrow & Y(fa,fb) \end{array} \quad \begin{array}{ccc} S^n & \longrightarrow & (a, X, b) \\ \downarrow & & \downarrow \\ D^{n+1} & \longrightarrow & (fa, Y, fb) \end{array}$$

is the same as giving a commutative square in **Top**, as on the right in the previous display, by $\sigma \dashv h$. The square on the right admits a diagonal filler $D^{n+1} \rightarrow X$ since f is a trivial $\mathcal{J}$ -fibration, and thus so does the square on the left.

(2): We proceed by induction on n . Having the right lifting property with respect to the inclusions

$$\emptyset \hookrightarrow 1 \quad 1+1 = \partial I \hookrightarrow I$$

ensures that f surjective and injective on path components, and thus is inverted by π_0 . For the inductive step we assume that $U_n \pi_n$ sends trivial $\mathcal{J}$ -fibrations to trivial $\mathcal{I}_{\leq n}$ -fibrations and suppose that f is a trivial $\mathcal{J}$ -fibration. Then so are all the maps it induces between path spaces by (1). But from the inductive definition of Trm_{n+1} we have $U_{n+1} \pi_{n+1} = \mathcal{G}(u_n \pi_n)P$ and so $U_{n+1} \pi_{n+1}(f)$ is a morphism of $(n+1)$ -globular sets which is surjective on objects (as argued already in the $n = 0$ case) and whose hom maps are trivial $\mathcal{I}_{\leq n}$ -fibrations by induction. Thus the result follows by lemma(56). $\square$

Corollary 61. ([11] Theorem(4.8)) *Trm_n is the category of algebras for a contractible $\mathcal{T}_{\leq n}$ -operad.*

PROOF. We proceed by induction on n , and the case $n = 0$ holds trivially. For the inductive step we must show by corollary(59) that the components (9) are trivial $\mathcal{I}_{\leq n}$ -fibrations, where $Q = \pi_n$, $T = \mathcal{T}_{\leq n}$ and $\phi : S \rightarrow T$ the contractible operad for Trimble n -categories. But by lemma(60) the unique map $U^S Q(A) \rightarrow 1$ is a trivial $\mathcal{I}_{\leq n}$ -fibration since A is contractible, and so the result follows from lemma(54) since trivial fibrations are closed under products and composition. $\square$

Acknowledgements

The author would like to acknowledge Michael Batanin, Clemens Berger, Denis-Charles Cisinski and Paul-André Melliès for interesting discussions on the substance of this paper. He would also like to express his gratitude to the laboratory PPS (Preuves Programmes Systèmes) in Paris, the Max Planck Institute in Bonn, the IHES and the Macquarie University Mathematics Department for the excellent working conditions he enjoyed during this project.

References

- [1] J. Adamek, F. Borceux, S. Lack, and J. Rosicky. A classification of accessible categories. *JPAA*, 175:7–30, 2002.
- [2] J. Adamek and J. Rosicky. *Locally presentable and accessible categories*. Number 189 in London Math Soc. Lecture Notes. Cambridge University Press, 1994.
- [3] M. Batanin. Monoidal globular categories as a natural environment for the theory of weak n -categories. *Advances in Mathematics*, 136:39–103, 1998.
- [4] M. Batanin, D-C. Cisinski, and M. Weber. Algebras of higher operads as enriched categories II. Available at <http://sites.google.com/site/markwebersmaths/>.
- [5] M. Batanin, D-C. Cisinski, and M. Weber. The lifting theorem for multitensors. Available at <http://sites.google.com/site/markwebersmaths/>.
- [6] M. Batanin and M. Weber. Algebras of higher operads as enriched categories. To appear in *Applied Categorical Structures* available at <http://www.pps.jussieu.fr/~weber/>, 2008.
- [7] J. Beck. Distributive laws. *Lecture Notes in Math.*, 80:119–140, 1969.
- [8] C. Berger, P-A. Melliès, and M. Weber. Monads with arities and their associated theories. Available at <http://sites.google.com/site/markwebersmaths/>.
- [9] A. Carboni, S. Lack, and R.F.C Walters. Introduction to extensive and distributive categories. *JPAA*, 84:145–158, 1993.

- [10] E. Cheng. Iterated distributive laws. arXiv:0710.1120v1, 2007.
- [11] E. Cheng. Comparing operadic theories of n -category. arXiv:0809.2070v2, 2008.
- [12] J.R.B. Cockett. Introduction to distributive categories. *Mathematical Structures in Computer Science*, pages 277–307, 1993.
- [13] S. Crans. A tensor product for Gray categories. *Theory and applications of categories*, 5:12–69, 1999.
- [14] B. Day and R. Street. Lax monoids, pseudo-operads, and convolution. In *Diagrammatic Morphisms and Applications*, volume 318 of *Contemporary Mathematics*, pages 75–96, 2003.
- [15] P. Gabriel and F. Ulmer. *Lokal Praesentierbare Kategorien*. Number 221 in SLNM. Springer Verlag, 1971.
- [16] R. Gordon, A.J. Power, and R. Street. *Coherence for tricategories*. Number 117 in Mem. Amer. Math. Soc. AMS, 1995.
- [17] G.M. Kelly. *Basic concepts of enriched category theory, LMS lecture note series*, volume 64. Cambridge University Press, 1982.
- [18] G.M. Kelly and S. Lack. V -Cat is locally presentable or locally bounded if V is so. *Theory and applications of categories*, 8:555–575, 2001.
- [19] S. Lack. Codescent objects and coherence. *J. Pure Appl. Algebra*, 175:223–241, 2002.
- [20] S. Lack and R.H. Street. The formal theory of monads II. *J. Pure Appl. Algebra*, 175:243–265, 2002.
- [21] T. Leinster. *Higher operads, higher categories*. Lecture note series. London Mathematical Society, 2003.
- [22] M. Makkai and R. Paré. *Accessible Categories*, volume 104 of *Contemp. Math.* AMS, 1989.
- [23] P-A. Melliès. Segal condition meets computational effects. <http://www.pps.jussieu.fr/mellies/papers/segal-lics-2010.pdf>.
- [24] R. Street. The formal theory of monads. *J. Pure Appl. Algebra*, 2:149–168, 1972.
- [25] R. Street. The role of Michael Batanin’s monoidal globular categories. *Contemporary Mathematics*, 230:99–116, 1998.
- [26] M. Weber. Familial 2-functors and parametric right adjoints. *Theory and applications of categories*, 18:665–732, 2007.
- [27] M. Weber. Free products of higher operad algebras. Available at <http://sites.google.com/site/markwebersmaths/>, 2009.

DEPARTMENT OF MATHEMATICS, MACQUARIE UNIVERSITY
 E-mail address: mark.weber.math@gmail.com